















- [8] Yatsuya H, Li Y, Hilawe EH, Ota A, Wang C, Chiang C, et al. Global trend in overweight and obesity and its association with cardiovascular disease incidence. Vol. 78, *Circulation Journal. Japanese Circulation Society*; 2014. p. 2807–18.
- [9] Kelly T, Yang W, Chen CS, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes*. 2008 Sep;32(9):1431–7.
- [10] Anzaku AS. Prevalence and Risk Factors for Obesity in a Nigerian Obstetric Population. *Am J Heal Res*. 2014;2(5):229.
- [11] Anzaku A, Idikwu O, Emmanuel O, Kingsley O. Impacts of Obesity on Maternal and Fetal Outcomes in Women with Singleton Pregnancy at a Nigerian Clinical Setting. *Br J Med Med Res*. 2015;6(12):1159–65.
- [12] Chigbu C, Aja L. Obesity in pregnancy in southeast Nigeria. *Ann Med Health Sci Res*. 2011;1(2):135–13540.
- [13] Ezeanochie MC, Ande AB, Olagbuji BN. Maternal obesity in early pregnancy and subsequent pregnancy outcome in a Nigerian population. *Afr J Reprod Health*. 2011;15(4):55–9.
- [14] Ramsey, P. S Schenken RS. Obesity in pregnancy: Complications and maternal management. Up To Date. 2019;1–40.
- [15] Okafor II. Indications for Caesarean Delivery in a State University Teaching Hospital, Enugu, Southeast, Nigeria. *Int J Nurs Healthc*. 2015;1(1):16–21.
- [16] Betrán AP, Ye J, Moller A-B, Zhang J, Gülmezoglu AM, Torloni MR. The Increasing Trend in Caesarean Section Rates: Global, Regional and National Estimates: 1990-2014. *PLoS One*. 2016;11(2): e0148343.
- [17] Betran AP, Torloni MR, Zhang JJ, Gülmezoglu A, Zongo A. WHO Statement on Caesarean Section Rates. *BJOG*. 2016; 123:667–70.
- [18] Tipton AM, Cohen SA, Chelmow D. Wound infection in the obese pregnant woman. Vol. 35, *Seminars in Perinatology*. W.B. Saunders; 2011. p. 345–9.
- [19] Kadia BM, Chichom-Mefire A, Halle-Ekane GE. Exploring the role of obesity and overweight in predicting the post-operative outcome of abdominal surgery in a sub-Saharan African setting: a prospective cohort study. *BMC Res Notes*. 2018 Dec;11(1):742.
- [20] Thornburg LL. Antepartum obstetrical complications associated with obesity. Vol. 35, *Seminars in Perinatology*. 2011. p. 317–23.
- [21] Tan T, Sia AT. Anaesthesia considerations in the obese gravida. Vol. 35, *Seminars in Perinatology*. 2011. p. 350–5.
- [22] Cleveland RD, Zitsch RP, Laws HL. Incisional hernia in morbidly obese patients. *The American Surgeon*. 1989; 55(1): 61-63.
- [23] Sauerland S, Korenkov M, Kleinen T, Arndt M, Paul A. Obesity is a risk factor for recurrence after incisional hernia repair. *Hernia*. 2004; 8(1): 42-46.
- [24] Gunatilake RP, Perlow JH. Obesity and pregnancy: Clinical management of the obese gravida. Vol. 204, *American Journal of Obstetrics and Gynecology*. 2011. p. 106–19.
- [25] WALL P. Vertical skin incisions and wound complications in the obese parturient. *Obstet Gynecol*. 2003 Nov;102(5):952–6.
- [26] Ceydeli A, Rucinski J, Wise L. Finding the best abdominal closure: An evidence-based review of the literature. *Curr Surg*. 2005;62(2):220–5.
- [27] Walton RB, Shnaekel KL, Ounpraseuth ST, Napolitano PG, Magann EF. High transverse skin incisions may reduce wound complications in obese women having cesarean sections: a pilot study. *J Matern Neonatal Med*. 2019 Mar;32(5):781–5.
- [28] Aabakke AJM, Krebs L, Ladelund S, Secher NJ. Incidence of incisional hernia after cesarean delivery: A register-based cohort study. *PLoS One*. 2014 Sep;9(9).
- [29] Ayandipo OO, Afuwape OO, Irabor DO, Abdurrazzaaq AI. Adult Abdominal Wall Hernia in Ibadan. *Ann Ibadan Postgrad Med [Internet]*. 2015;13(2):94–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27162521>0 A & <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4853882>.