



























Med. 2019;20(S1):3–11.

[21] Pry J, Chipungu J, Smith HJ, Bolton Moore C, Mutale J, Duran-Frigola M, et al. Patient-reported reasons for declining same-day antiretroviral therapy initiation in routine HIV care settings in Lusaka, Zambia: results from a mixed-effects regression analysis. *J Int AIDS Soc.* 2020 Jul;23(7):e25560.

[22] Rufu A, Chitimbire VTS, Nzou C, Timire C, Owiti P, Harries AD, et al. Implementation of the “Test and Treat” policy for newly diagnosed people living with HIV in Zimbabwe in 2017. *Public Health Action.* 2018 Sep 21;8(3):145–50.

[23] Factors Influencing Rapid Antiretroviral Therapy Initiation at Four eThekweni Clinics, KwaZulu-Natal, South Africa | SpringerLink [Internet]. [cited 2021 Dec 15]. Available from: <https://link.springer.com/article/10.1007/s10461-021-03530-3>.

[24] Ahmed I, Demissie M, Worku A, Gugsu S, Berhane Y. Effectiveness of same-day antiretroviral therapy initiation in retention outcomes among people living with human immunodeficiency virus in Ethiopia: empirical evidence. *BMC Public Health.* 2020 Nov 26;20(1):1802.

[25] Katbi M, Adegboye A, Adedoyin A, Yunusa F, Kayode G, Bello M, et al. Effect of clients Strategic Index Case Testing on community-based detection of HIV infections (STRICT study). *Int J Infect Dis IJID Off Publ Int Soc Infect Dis.* 2018 Sep;74:54–60.

[26] Rosen S, Maskew M, Fox MP, Nyoni C, Mongwenyana C, Malete G, et al. Initiating Antiretroviral Therapy for HIV at a Patient’s First Clinic Visit: The RapIT Randomized Controlled Trial. *PLoS Med.* 2016 May;13(5):e1002015.