

The Impact of Dietary Cholesterol on LDL Cholesterol Levels in Adults: A Short-Term Observational Study

Iman A. Elbanna^{1*}, Naveen Karthik P², Thangaswamy Selvankumar³

¹*Urgent System of Care Lead, Model of care, Ministry of Health, Hail region, Saudi Arabia*

²*Department of Orthopaedics, Saveetha Medical College & Hospital, Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai-602105, Tamil Nadu, India*

³*Department of General Medicine, Saveetha Medical College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, 602105, Tamil Nadu, India*

Abstract

Dietary cholesterol has long been implicated in regulating blood cholesterol levels, particularly low-density lipoprotein (LDL) cholesterol, a key marker of cardiovascular health. Understanding the role of dietary cholesterol is critical, given its potential influence on the development of cardiovascular diseases (CVD). This short-term observational study evaluated the effects of high-cholesterol and low-cholesterol diets on LDL cholesterol levels in healthy adults aged 25-50 years. A total of 160 participants were randomly assigned to either a high-cholesterol or low-cholesterol diet for six months. Dietary intake was monitored using food frequency questionnaires (FFQs) and self-reported food diaries while fasting blood samples were collected at baseline, three months, and six months to measure LDL cholesterol, total cholesterol, HDL cholesterol, and triglycerides. Results revealed a modest 8.3% increase in LDL cholesterol in the high-cholesterol group ($p < 0.05$), while the low-cholesterol group exhibited a slight but statistically insignificant decrease of 3.4% ($p > 0.05$). Other lipid parameters, such as HDL cholesterol and triglycerides, remained stable in both groups. The findings suggest that dietary cholesterol may have a limited but measurable impact on LDL cholesterol levels, without significantly influencing overall cardiovascular risk in the short term. These results challenge traditional views on dietary cholesterol and emphasize the importance of individualized dietary recommendations. This study contributes valuable insights to the ongoing discussion surrounding cholesterol management and dietary guidelines.

Keywords: Cardiovascular Risk, Dietary Cholesterol, LDL Cholesterol, Lipid Metabolism, Lipid Parameters.

Introduction

Cholesterol is an essential lipid in the human body, playing a vital role in maintaining cellular integrity, synthesizing steroid hormones, and producing vitamin D [1, 2]. It is predominantly synthesized by the liver and obtained through dietary sources [3, 4], with animal-based foods such as eggs, meat, and shellfish being primary contributors [5, 6]. While cholesterol is indispensable for

physiological functions, elevated levels of low-density lipoprotein (LDL) cholesterol are strongly associated with an increased risk of cardiovascular diseases (CVD), particularly atherosclerosis, a condition characterised by plaque buildup in arterial walls [7, 8]. For decades, dietary cholesterol has been viewed as a key contributor to elevated LDL cholesterol levels and associated cardiovascular risk. This perception led to dietary guidelines recommending stringent limitations on

cholesterol-rich foods [9, 10]. However, recent research suggests that the body's endogenous cholesterol synthesis adjusts in response to dietary intake, potentially minimizing the impact of dietary cholesterol on blood lipid levels in most individuals [11, 12]. These findings challenge conventional dietary recommendations and highlight the complexity of cholesterol metabolism.

Despite growing evidence, the precise relationship between dietary cholesterol and LDL cholesterol remains a topic of debate. Some studies suggest that certain individuals, known as "hyper-responders," may exhibit significant changes in LDL cholesterol levels in response to increased dietary cholesterol, while others experience minimal or negligible effects [9, 10]. Additionally, confounding factors such as baseline dietary habits, physical activity, and genetic predispositions further complicate this relationship [13, 14]. This study aims to address these uncertainties by assessing the short-term effects of high-cholesterol and low-cholesterol diets on LDL cholesterol levels in healthy adults aged 25-50 years. By incorporating a randomized design and accounting for potential confounding factors, this research seeks to provide a clearer understanding of how dietary cholesterol influences lipid profiles. Furthermore, the study explores the implications of these findings for public health and dietary guidelines, with the ultimate goal of informing more nuanced and individualized approaches to cholesterol management.

The primary objective of this study was to assess the short-term effects of dietary cholesterol intake on LDL cholesterol levels in healthy adults aged 25-50 years. Specifically, the study aimed to examine the impact of increased dietary cholesterol intake on LDL cholesterol levels over six months. Investigate whether a low-cholesterol diet could result in measurable reductions in LDL cholesterol levels. Evaluate changes in additional lipid parameters, including total cholesterol, high-density lipoprotein (HDL) cholesterol, and

triglycerides, in response to varying dietary cholesterol levels. Identify potential influences of confounding factors, such as physical activity, smoking, alcohol consumption, and baseline dietary habits, on lipid profile outcomes. Explore the role of individual variability in cholesterol metabolism, including the differences observed between hyper-responders and hypo-responders.

Materials and Methods

Study Design

This short-term, 6-month observational study employed a randomized design to assess the impact of dietary cholesterol on LDL cholesterol levels. Participants were randomly assigned to a high-cholesterol diet group or a low-cholesterol diet group, and their lipid profiles were monitored at baseline, 3 months, and 6 months [15].

Participants

A total of 160 healthy adults aged 25-50 years were recruited. The participants were stratified by gender, ethnicity, and baseline dietary habits to ensure a representative sample [16].

Inclusion Criteria

1. Healthy adults aged 25-50 years.
2. No history of cardiovascular diseases, diabetes, or hyperlipidemia.
3. Non-smokers or those with stable smoking habits.
4. No use of cholesterol-lowering medications.

Exclusion Criteria

1. Pregnant or breastfeeding women.
2. Individuals with medical conditions affecting lipid metabolism [17].

Diet Intervention

Participants were assigned to one of two groups:

1. High-Cholesterol Diet Group: Instructed to increase daily cholesterol intake to

approximately 300 mg/day through eggs, shellfish, and dairy products [18].

2. **Low-Cholesterol Diet Group:** Instructed to limit cholesterol intake to below 100 mg/day, focusing on plant-based foods and avoiding animal-based cholesterol sources [19].

Data Collection

1. **Dietary Monitoring:** Food diaries intake. Follow-up interviews ensured adherence to assigned diets [20].
2. **Blood Biomarkers:** Fasting blood samples were collected at baseline, 3 months, and 6 months to measure LDL cholesterol, total cholesterol, HDL cholesterol, and triglycerides [21, 22].
3. **Confounding Factors:** Physical activity, smoking, and alcohol consumption were assessed through questionnaires [23].
4. **Food frequency questionnaires (FFQs)** were used to monitor dietary.

Statistical Analysis

Data were analyzed using statistical software to compare lipid profile changes between the

two groups. Paired and unpaired t-tests were employed to evaluate within-group and between-group differences, respectively. Statistical significance was set at $p < 0.05$, and additional analyses were conducted to account for individual variability in cholesterol response [24].

This robust methodology ensured the reliability and validity of the study's findings, providing a solid foundation for understanding the short-term effects of dietary cholesterol on LDL cholesterol levels and other lipid parameters.

Results

The dietary cholesterol intake of the different groups was studied. Table 1 depicts the LDL cholesterol levels for the 3 months of the study. From the observations the obtained results are

1. **High-Cholesterol Group:** Increased dietary cholesterol intake to 300 mg/day.
2. **Low-Cholesterol Group:** Reduced dietary cholesterol intake to <100 mg/day.

Table 1. LDL Cholesterol Levels

S.No.	Time Point	High-Cholesterol Group (mg/dl)	Low-Cholesterol Group (mg/dl)
1	Baseline	120	118
2	3 Months	125	116
3	6 Months	130	114

Table 2 describes the other lipid parameters of total cholesterol, HDL cholesterol and triglycerides. The significant results are

1. **High-Cholesterol Group:** LDL cholesterol increased by 8.3% ($p < 0.05$).
2. **Low-Cholesterol Group:** LDL cholesterol decreased by 3.4% ($p > 0.05$).

Table 2. Other Lipid Parameters

S.No.	Parameter	High-Cholesterol Group	Low-Cholesterol Group
1	Total Cholesterol	Slight increase	Slight decrease
2	HDL Cholesterol	Stable	Stable
3	Triglycerides	No significant change	No significant change

Table 3 represents the summary of lipid profile changes in the High-Cholesterol Group and Low-Cholesterol Group.

Table 3. Summary of Lipid Profile Changes

S.No.	Lipid Parameter	High-Cholesterol Group	Low-Cholesterol Group
1	LDL Cholesterol (Δ)	+8.3%	-3.4%
2	HDL Cholesterol (Δ)	No change	No change
3	Total Cholesterol (Δ)	Slight increase	Slight decrease
4	Triglycerides (Δ)	No change	No change

The Change in LDL Cholesterol Levels over 6 Months is given in the Figure 1. It indicates LDL Cholesterol, HDL Cholesterol, Total

Cholesterol and Triglycerides changes in the High and low Cholesterol diet groups.




Figure 1. Change in LDL Cholesterol Levels Over 6 Months

(Placeholder for Graph Showing the Trend of LDL Cholesterol in Both Groups.)

Discussion

The study demonstrated that dietary cholesterol intake led to a modest increase in LDL cholesterol levels in the high-cholesterol

group, while the low-cholesterol diet had minimal effects. These findings align with emerging research that suggests dietary cholesterol plays a smaller role in determining

blood cholesterol levels than previously believed [25].

Individual Variability in Cholesterol Metabolism

One key observation from the study was the variation in LDL cholesterol responses among participants. This variability could be attributed to differences in cholesterol metabolism, with certain individuals being classified as "hyper-responders." These individuals experience greater increases in LDL cholesterol when dietary cholesterol intake rises, likely due to genetic or metabolic factors. On the other hand, "hypo-responders" exhibit minimal changes, as their bodies are more effective at compensating by reducing endogenous cholesterol synthesis. Future research should prioritize identifying biomarkers or genetic markers that can predict an individual's response to dietary cholesterol [26, 27].

Dietary Compensation and Homeostasis

Another critical factor is the body's ability to regulate cholesterol levels through dietary compensation. This physiological mechanism may explain why the observed LDL cholesterol increase in the high-cholesterol group was modest, even though dietary cholesterol intake was significantly elevated. The liver, which plays a central role in cholesterol synthesis and metabolism, likely adjusts production levels to maintain homeostasis. This finding underscores the complexity of lipid metabolism and highlights the importance of considering these regulatory processes in dietary recommendations.

Comparison with Previous Research

The results of this study are consistent with findings from Lichtenstein and Deckelbaum (2001), who reported that dietary cholesterol has a limited impact on LDL levels for most individuals. Similarly, Reddy and Teff (2012) suggested that other dietary components, such as saturated fat, may significantly influence LDL cholesterol more than dietary cholesterol

alone. While this study was limited to a 6-month duration, the overall trends align with long-term observational data suggesting that dietary cholesterol has a less pronounced effect on cardiovascular risk than previously assumed.

Role of Confounding Factors

Despite controlling for physical activity, smoking, and baseline dietary habits, these variables may still have influenced the study's outcomes. Physical activity, for instance, can independently lower LDL cholesterol levels and may have attenuated the effects of dietary changes in both groups. Similarly, variations in baseline dietary patterns, including differences in fibre or saturated fat intake, could have moderated the impact of cholesterol consumption. These factors highlight the need for comprehensive assessments in future studies.

Public Health Implications

The findings from this study challenge traditional dietary guidelines that emphasize stringent cholesterol restrictions. Instead, they support a shift toward more holistic dietary patterns that consider overall nutrient balance rather than isolating specific components. For example, the Mediterranean diet [28, 29], which includes moderate cholesterol intake but emphasizes fruits, vegetables, whole grains, and healthy fats, has consistently been associated with improved cardiovascular outcomes. Educating the public on the relative insignificance of dietary cholesterol for most individuals could reduce unnecessary dietary restrictions and promote more sustainable eating habits [30].

Limitations of the Study

While the findings are insightful, several limitations must be acknowledged. The short duration of the study (6 months) may not capture the long-term effects of dietary cholesterol on LDL levels or cardiovascular risk. Additionally, reliance on self-reported

dietary data introduces potential recall bias, which could affect the accuracy of dietary intake measurements. Finally, the study's design did not explore the impact of other dietary components, such as fibre or saturated fat, which may interact with cholesterol metabolism. Future research should address these limitations by employing longer study durations, objective dietary assessment methods, and more comprehensive analyses of dietary factors.

Future Research Directions

Building on these findings, future studies should:

1. Investigate the long-term effects of dietary cholesterol on lipid profiles and cardiovascular outcomes.
2. Include a more diverse population sample to explore demographic variations in cholesterol metabolism.
3. Examine interactions between dietary cholesterol and other nutrients, such as saturated fats and dietary fibre.
4. Utilize advanced lipid profiling techniques to assess changes in LDL particle size and composition, which may provide deeper insights into cardiovascular risk.

In conclusion, while dietary cholesterol can modestly affect LDL cholesterol levels in certain individuals, its overall impact appears to be limited for the general population. These findings contribute to a growing body of evidence that calls for a more nuanced approach to dietary guidelines, emphasizing overall dietary patterns rather than isolated nutrient restrictions.

Limitations

1. Short duration (6 months) may not capture long-term effects.
2. Reliance on self-reported dietary data introduces potential recall bias.
3. Individual variability in cholesterol metabolism was not fully explored.

Conclusion

The findings of this study underscore the nuanced relationship between dietary cholesterol intake and LDL cholesterol levels in healthy adults. While the high-cholesterol diet group exhibited a statistically significant increase in LDL cholesterol levels, the changes were modest and did not suggest immediate cardiovascular risk for most participants. Conversely, the low-cholesterol diet group demonstrated only minimal reductions in LDL cholesterol, reinforcing the idea that dietary cholesterol may not be a primary determinant of lipid profiles for the general population.

These results challenge long-standing dietary guidelines that advocate for strict avoidance of cholesterol-rich foods and instead highlight the importance of personalized dietary recommendations. The study's observations of individual variability in cholesterol metabolism, particularly among hyper-responders, suggest that a one-size-fits-all approach may not be appropriate for cholesterol management. Moreover, the findings emphasize the need for balanced diets that consider overall nutrient interactions rather than isolating specific components like cholesterol.

Future research should prioritize long-term investigations with diverse populations to explore the interplay between dietary cholesterol, genetic factors, and other dietary components such as saturated fats and fibre. Additionally, advanced lipid profiling techniques could offer more comprehensive insights into the cardiovascular implications of dietary cholesterol. Ultimately, this study contributes valuable data to the evolving discourse on cholesterol management [31–33], paving the way for more nuanced public health guidelines.

Emphasize Balanced Eating Patterns

Encourage individuals to adopt dietary patterns such as the Mediterranean or DASH diets, which focus on whole grains, fruits,

vegetables, lean proteins, and healthy fats. These patterns have consistently been shown to improve lipid profiles and reduce cardiovascular risk.

1. Moderation in Dietary Cholesterol:

Public health messaging should focus on moderate consumption of cholesterol-rich foods rather than outright avoidance. For most individuals, dietary cholesterol has minimal impact on LDL cholesterol levels when consumed as part of a balanced diet.

2. Nutrient Synergy: Highlight the importance of considering interactions between dietary cholesterol and other nutrients, such as saturated fats and dietary fibre, which play significant roles in cholesterol metabolism and overall cardiovascular health.

3. Tailored Recommendations: Develop dietary guidelines that account for individual variability in cholesterol response [34]. For instance, those identified as "hyper-responders" may benefit from specific advice to limit cholesterol-rich foods.

Public Health Policy

1. Education Campaigns: Launch public awareness initiatives to dispel myths surrounding dietary cholesterol and its impact on heart health. Focus on promoting evidence-based dietary patterns rather than demonizing individual nutrients.

2. Nutritional Labeling: Update food labeling standards to provide clear, actionable information about the overall nutrient profile of foods, rather than focusing solely on cholesterol content. This approach can help consumers make informed choices.

3. Supportive Environments: Encourage the food industry to reformulate products by reducing saturated fats and enhancing the availability of high-fibre and nutrient-dense options that support heart health.

4. Equity in Dietary Recommendations:

Address socioeconomic and cultural barriers to healthy eating by ensuring affordable and accessible options for balanced diets across diverse populations.

Clinical Practice

1. Routine Lipid Monitoring: Clinicians should monitor lipid profiles periodically, especially for individuals at higher risk of cardiovascular disease or those with significant dietary changes.

2. Personalized Nutrition Counseling: Encourage healthcare providers to adopt a personalized approach in dietary counselling, recognizing that responses to dietary cholesterol vary widely among individuals.

3. Incorporate Emerging Evidence: Update clinical guidelines to reflect current research on the limited role of dietary cholesterol in influencing LDL cholesterol for the majority of the population.

Future Research

1. Long-Term Studies: Support research that examines the long-term effects of dietary cholesterol on lipid profiles and cardiovascular outcomes, including studies with larger and more diverse populations.

2. Advanced Biomarkers: Explore the use of advanced lipid profiling techniques to understand how dietary cholesterol influences LDL particle size, density, and other cardiovascular risk factors.

3. Genetic and Metabolic Factors: Investigate genetic markers and metabolic pathways that determine individual variability in cholesterol response, paving the way for precision nutrition strategies.

4. Interactions with Other Dietary Components: Conduct studies on the interactions between dietary cholesterol and other nutrients, such as dietary fibre and polyunsaturated fats [20], to develop comprehensive dietary recommendations.

Lifestyle Integration

1. **Physical Activity:** Reinforce the importance of regular physical activity in conjunction with dietary interventions for optimal cardiovascular health.
2. **Behavioral Strategies:** Develop and implement behavior-change interventions to help individuals sustain healthy eating patterns over the long term.
3. **Community Support:** Encourage community-based programs that provide education, resources, and support for adopting heart-healthy dietary habits.

These expanded recommendations aim to provide actionable, evidence-based strategies to promote heart health while addressing the complexities of dietary cholesterol and its impact on lipid metabolism.

Acknowledgement

The authors would like to express their gratitude to all participants who generously

gave their time and effort to contribute to this study. We also acknowledge the support provided by the Ministry of Health, Hail Region, Saudi Arabia, and the Department of General Medicine at Saveetha Medical College and Hospital, Saveetha University, Chennai, India, for their guidance and resources throughout the research process. Special thanks to the laboratory staff and data collection team for their meticulous work in ensuring the accuracy and reliability of the data. Lastly, we extend our appreciation to our families and colleagues for their encouragement and support.

Conflict of Interest

The authors declare no conflicts of interest regarding the publication of this article. The research was conducted independently, without any influence from commercial entities, sponsors, or funding agencies that could have biased the study's design, results, or conclusions.

References

- [1]. Sharma, B., Agnihotri, N., 2019. Role of cholesterol homeostasis and its efflux pathways in cancer progression. *J Steroid Biochem Mol Biol.* 191: 105377. doi:10.1016/j.jsbmb.2019.105377
- [2]. Carson, J. A. S., Lichtenstein, A. H., Anderson, C. A. M., Appel, L. J., Kris-Etherton, P. M., Meyer, K. A., et al., 2020., Dietary cholesterol and cardiovascular risk: A science advisory from the American heart association. *Circulation.* 141: e39–e53. doi:10.1161/CIR.0000000000000743
- [3]. Schwab, U., Reynolds, A. N., Sallinen, T., Rivelles, A. A., Risérus, U., 2021., Dietary fat intakes and cardiovascular disease risk in adults with type 2 diabetes: a systematic review and meta-analysis. *Eur J Nutr.* 60: 3355–3363. doi:10.1007/s00394-021-02507-1
- [4]. Stellaard, F., 2022., From dietary cholesterol to blood cholesterol, physiological lipid fluxes, and cholesterol homeostasis. *Nutrients.* 14: 1643. doi:10.3390/nu14081643

- [5]. Shahoei, S. H., Nelson, E. R., 2019., Nuclear receptors, cholesterol homeostasis and the immune system. *J Steroid Biochem Mol Biol.* 191: 105364. doi:10.1016/j.jsbmb.2019.04.013
- [6]. Szczepańska, E., Białek-Dratwa, A., Janota, B., Kowalski, O., 2022., Dietary therapy in prevention of cardiovascular disease (CVD)—tradition or modernity? A review of the latest approaches to nutrition in CVD. *Nutrients.* 14. doi:10.3390/nu14132649
- [7]. Willett, W., 2017., Eat, drink, and be healthy: The Harvard medical school guide to healthy eating. Free Press; Available: <https://books.google.com/books?hl=en&lr=&id=uk1DwAAQBAJ&oi=fnd&pg=PP7&dq=Eat,+Drink,+and+Be+Healthy:+The+Harvard+Medical+School+Guide+to+Healthy+Eating.+Free+Press&ots=6IPwSAIqF3&sig=w37mVc9gKLD8trkI41YzpoLZtWs>
- [8]. Soliman, G. A., 2018., Dietary cholesterol and the lack of evidence in cardiovascular disease. *Nutrients.* 10: 780. doi:10.3390/nu10060780

- [9]. Schade, D. S., Shey, L., Eaton, R. P., 2020., Cholesterol review: a metabolically important molecule. *Endocr Pract.* Available: <https://www.sciencedirect.com/science/article/pii/S1530891X20484472>
- [10]. Sacks, F. M., Lichtenstein, A. Ho., Wu, J. H. .Y., Appel, L. J., Creager, M. A., Kris-Etherton, P. M., et al., 2017., Dietary fats and cardiovascular disease: A presidential advisory from the American Heart Association. *Circulation.* 136: e1–e23. doi:10.1161/CIR.0000000000000510
- [11]. Zhu, Y., Bo, Y., Liu, Y., 2019., Dietary total fat, fatty acids intake, and risk of cardiovascular disease: a dose-response meta-analysis of cohort studies. *Lipids Health Dis.* 18: 91. doi:10.1186/s12944-019-1035-2
- [12]. Minervini, G., Franco, R., Marrapodi, M.M., Di Blasio. M., Ronsivalle, V., Cicciù, M., 2023., Children oral health and parents education status: a cross-sectional study. *BMC Oral Health.* 24; 23(1):787. doi: 10.1186/s12903-023-03424-x.
- [13]. Yanai, H., Tada, N., 2018., Effects of consumption of various fatty acids on serum HDL-cholesterol levels. *Journal of Endocrinology and Metabolism.* 8: 94–99. doi:10.14740/JEM.V8I5.534
- [14]. Jenkins, W., Jenkins, A., Jenkins, A., Brydson, C., 2019., The portfolio diet for cardiovascular disease risk reduction: an evidence based approach to lower cholesterol through plant food consumption. Available: https://books.google.com/books?hl=en&lr=&id=S_tUDwAAQBAJ&oi=fnd&pg=PP1&dq=The+portfolio+diet+for+cardiovascular+risk+reduction:+An+evidence-based+approach+to+lower+cholesterol+levels&ots=pZDea6M_Dk&sig=1HdK9OsiUSrEhoddKORogXBdFuc
- [15]. Doma, K. M., Dolinar, K. F., Dan Ramdath, D., Wolever, T. M. S., Duncan, A. M., 2021., Canned beans decrease serum total and LDL cholesterol in adults with elevated LDL cholesterol in a 4-wk multicenter, randomized, crossover study. *J Nutr.* 151: 3701–3709. doi:10.1093/jn/nxab323
- [16]. Lordan, R., Tsoupras, A., Mitra, B., Zabetakis, I., 2018., Dairy fats and cardiovascular disease: Do we really need to be concerned? *Foods.* 7: 29. doi:10.3390/foods7030029
- [17]. Yu, E., Malik, V. S., Hu, F. B., 2018., Cardiovascular disease prevention by diet modification: JACC health promotion series. *J Am Coll Cardiol.* 72: 914–926. doi:10.1016/j.jacc.2018.02.085
- [18]. Lawrence, G. D., 2021., Perspective: The saturated fat-unsaturated oil dilemma: Relations of dietary fatty acids and serum cholesterol, atherosclerosis, inflammation, cancer, and all-cause mortality. *Adv Nutr.* ;12: 647–656. doi:10.1093/advances/nmab013
- [19]. Drouin-Chartier, J-P., Chen, S., Li, Y., Schwab, A. L., Stampfer, M. J., Sacks, F. M., et al., 2020., Egg consumption and risk of cardiovascular disease: three large prospective US cohort studies, systematic review, and updated meta-analysis. *BMJ.* 368: m513. doi:10.1136/bmj.m513
- [20]. Trautwein, E. A., McKay, S., 2020., The role of specific components of a plant-based diet in management of dyslipidemia and the impact on cardiovascular risk. *Nutrients.* 12: 2671. doi:10.3390/nu12092671
- [21]. Ramadoss, R., Padmanaban, R., Subramanian, B., 2021., Role of bioglass in enamel remineralization: Existing strategies and future prospects—A narrative review. *Journal of Biomedical Materials Research Part B: Applied Biomaterials,* 110(1), 45-66. <https://doi.org/10.1002/jbm.b.34904>
- [22]. Gjuladin-Hellon, T., Davies, I. G., Penson, P., Amiri Baghbadorani, R., 2019., Effects of carbohydrate-restricted diets on low-density lipoprotein cholesterol levels in overweight and obese adults: a systematic review and meta-analysis. *Nutr Rev.* 77: 161–180. doi:10.1093/nutrit/nuy049
- [23]. Lloyd-Jones, D. M., Allen, N. B., Anderson, C. A. M., Black, T., Brewer, L. C., Foraker, R. E., et al., 2022., Life's Essential 8: Updating and enhancing the American heart association's construct of cardiovascular health: A presidential advisory from the American Heart Association. *Circulation.* 146: e18–e43. doi:10.1161/CIR.0000000000001078

- [24]. Dibaba, D. T., 2019., Effect of vitamin D supplementation on serum lipid profiles: a systematic review and meta-analysis. *Nutr Rev.* doi:10.1093/nutrit/nuz037
- [25]. Hooper, L., Martin, N., Jimoh, OF., Kirk, C., Foster, E., Abdelhamid, A. S., 2020., Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database Syst Rev.* 8: CD011737. doi:10.1002/14651858.CD011737.pub3
- [26]. Aguilar-Ballester, M., Herrero-Cervera, A., Vinué, Á., Martínez-Hervás, S., González-Navarro, H., 2020., Impact of cholesterol metabolism in immune cell function and atherosclerosis. *Nutrients.* 12. doi:10.3390/nu12072021
- [27]. Maki, K. C., Dicklin, M. R., Kirkpatrick, C. F., 2021., Saturated fats and cardiovascular health: Current evidence and controversies. *J Clin Lipidol.* 15: 765–772. doi:10.1016/j.jacl.2021.09.049
- [28]. Rashid, MF., Karobari MI, Halim, M.S., Noorani, T.Y., 2022., Effectiveness of Visual-Tactile Examination and DIAGNOdent Pen in Detecting Early Enamel Caries and Its Remineralisation: An In Vitro Study. *Biomed Res Int.* 11:1263750. doi: 10.1155/2022/1263750.
- [29]. Bouchard, J., Valookaran, A. F., Aloud, B. M., Raj, P., Malunga, L. N., Thandapilly, S. J., et al., 2022., Impact of oats in the prevention/management of hypertension. *Food Chem.* 381: 132198. doi:10.1016/j.foodchem.2022.132198
- [30]. Sellem, L., Flourakis, M., Jackson, K. G., Joris, P. J., Lumley, J., Lohner, S., et al., 2022., Impact of replacement of individual dietary SFAs on circulating lipids and other biomarkers of cardiometabolic health: A systematic review and meta-analysis of randomized controlled trials in humans. *Adv Nutr.* 13: 1200–1225. doi:10.1093/advances/nmab143
- [31]. DiNicolantonio, J. J., O’Keefe, J. H., 2018., Effects of dietary fats on blood lipids: a review of direct comparison trials. *Open Heart.* 5: e000871. doi:10.1136/openhrt-2018-000871
- [32]. Correction to: Dietary fats and cardiovascular disease: A presidential advisory from the American Heart Association. *Circulation.* 2017;136: e195. doi:10.1161/CIR.0000000000000529
- [33]. Pérez-Ramírez, I. F., Becerril-Ocampo, L. J., Reynoso-Camacho, R., Herrera, M. D., Guzmán-Maldonado, S. H., Cruz-Bravo, R. K., 2018., Cookies elaborated with oat and common bean flours improved serum markers in diabetic rats. *Journal of the Science of Food and Agriculture.* 98: 998–1007. Available: <http://zacatecas.inifap.gob.mx/publicaciones/perez-ramirez%20et%20al.%202018.pdf>
- [34]. Blesso, C., Fernández, M., 2018., Dietary cholesterol, serum lipids, and heart disease: Are eggs working for or against you? *Nutrients.* 10. doi:10.3390/nu10040426