

Determinants of Caregivers Knowledge Attitude and Practice Towards MCV2 in Adamawa State Nigeria, February 2024

Visa Ibrahim Tyakaray^{1*}, Martins Olutayo Folashade²

¹Department of Public Health, Texila American University, Guyana, South America

²Department of Public Health, Modibbo Adama University Teaching Hospital, Yola, Adamawa State, Nigeria

Abstract

Despite the availability of reliable and affordable measles vaccines, Nigeria is still experiencing low vaccination coverage, resulting in high cases of measles and related complication. This study assessed the determinants of caregivers' knowledge and attitude towards measles and MCV2 in Adamawa State, Nigeria. A three-arm, parallel randomized control trial was conducted in four health facilities sampled through three-stage multistage sampling technique in Adamawa State. Semi-structured interviewer-administered questionnaire was administered to 351 enrolled caregivers. Mean age of caregivers was 30.7 (SD±6.7); 212 (60.39%) of caregivers had good knowledge of measles and MCV2 and 265 (75.5%) had high level of positive attitude toward measles and MCV2. Major factors statistically associated with good knowledge of measles and MCV2 among caregivers include: age ($X^2=27.1$, $p\text{-value}=<0.001$) marital status ($X^2=37.0$, $p\text{-value}=<0.001$) and Family size ($X^2=15.1$, $p\text{-value}=<0.001$). Factors statistically associated with positive attitude of mothers towards MCV2 include: marital status ($X^2=49.0$, $p\text{-value}=<0.001$), religion ($X^2=73.3.0$, $p\text{-value}=<0.001$) and occupation of caregiver ($X^2=11.1$, $p\text{-value}=<0.049$) and factors associated with practice of timeliness towards MCV2 uptake were marital status, place of residence, and the age group of caregivers. Several factors influence caregivers' knowledge and attitude towards MCV2. Therefore, the Adamawa State government should implement measures to enhance measles vaccination coverage by increasing awareness of MCV2 through monitoring and addressing knowledge gaps on MCV2.

Keywords: Attitude, Caregivers, Determinants, Knowledge, MCV2.

Introduction

Immunization serves as the cornerstone of an effective health system, significantly contributing to the reduction of morbidity and mortality associated with various infectious diseases [1]. It is among the most successful health interventions and continues to be a vital means for achieving the Sustainable Development Goals (SDG) [2, 3].

The World Health Organization Immunization Agenda 2030 (IA2030) strives to reduce morbidity and mortality from vaccine-preventable diseases across the life course which is in consonant with the SDG 2030

priorities of eliminating vaccine preventable diseases and improving access to new lifesaving vaccines [4, 5]. However, for countries to attain the expected level of herd immunity, children have to receive the recommended doses of antigens. In Nigeria, the vaccination coverages of most antigens are still suboptimal with vaccine-preventable diseases accounting for an estimated 29% of childhood deaths annually [6].

The current immunization measles coverage rate falls short of national and global targets of 95% and 90% respectively [7, 8]. The global vaccination coverage survey (2010 to 2019)

showed decrease in first dose of Measles Containing Vaccine 1 (MCV1) between 84% and 86%, while measles second dose (MCV2) coverage increased from 42% to 71%, reflecting second dose introductions in many countries. From 2019 to 2020, global MCV1 coverage decreased to the 2014 level of 84%, whereas MCV2 coverage was relatively stable at 71% in 2019 and 70% in 2020. MCV1 coverage in 2020 ranged from 68% in the African Region [9]. In Adamawa State, nearly 1 in 4 of all the under-5 children in Adamawa State received no basic vaccinations [10].

Being a highly contagious disease that has caused global morbidity and mortality, tremendous progress has been made in measles elimination, yet resurgence of measles has been observed in recent years. [11]. Worldwide, more than 140,000 people died from measles in 2018, according to new estimates from the World Health Organization (WHO) and the United States Centers for Disease Control and Prevention (CDC) with most of the deaths occurring among children under the age of five years [12]. In Nigeria, a total of 203,089 measles cases were reported to the surveillance system with the overall incidence rate of 200 cases per 1 million population higher in the North exceeding that of the South of 24 cases per 1 million population yearly [13].

To reduce the burden of measles, the World Health Organization (WHO) recommended a coverage of 95% of 2 doses of measles-containing vaccine (MCV) to ensure an immunological response and stop transmission.

In order to interrupt measles transmission and prevent outbreaks, there is need to reach at least 95% coverage with both recommended doses of measles [14]. Studies have revealed that Nigeria is far from reaching its target due to suboptimal RI performance and high measles dropout rate which is a critical concern especially with the low vaccination coverage in northern Nigeria [15].

There is the paucity of information regarding the knowledge, attitude and practices of

caregivers towards MCV2 in Nigeria and Adamawa State in particular. Understanding the factors influencing mothers' knowledge and attitudes towards immunization could guide targeted interventions to improve MCV2 coverage. Therefore, this study was carried out to determine the determinants of caregiver's knowledge, attitudes, and practices towards MCV2 in Adamawa state.

Materials and Methods

Factors influencing caregivers' knowledge and attitudes were obtained from a three-arm, parallel, randomized controlled trial which is being conducted in four health facilities and involving 351 caregivers in Adamawa State Nigeria. The health facilities included Ngbalang & Gweda Mallam PHCCs of Numan LGA and Major Aminu & Jambutu PHCCs of Yola North LGA. Randomization was applied to health facilities during enrolment with a 1:1:1 allocation ratio of one control arm and two intervention arms.

Participant selection criteria and recruitment Criteria for inclusion into this study included all mothers and/or caregiver who had a 9-month-old child that is vaccinated with MCV1, aged 18 years and older, have a working mobile phone, have resided in the study area at least for 6 months before enrolment and willing to provide consent for participation in the study. Mothers who could not read mobile phone text messages in any of the four languages used in the study (Hausa, English languages, Fulfulde or Bwatiye), Mothers and/or caregivers who had no mobile network access in their house/compound and mothers who planned to relocate out of the study area during the study follow-up period.

The sample size was calculated using the formula by [16]. The largest sample size was obtained for the outcome variable knowledge on measles and MCV2. The sample frame was the list of patients at each recruitment site. Simple randomization with an allocation of 1:1:1 to the control, SMS or Automated Calls

were performed using a list of generated random numbers in all the selected health facilities (1 Control arm and 2 Intervention arms). Units of randomization were mother-infant pairs in each of the study arms of

interventions and control. Total number of caregivers assessed for eligibility were 507 caregivers; 156 caregivers were excluded based on the exclusion criteria outlined. (Figure 1).




Figure 1. Flow Chart of Randomized Controlled Trial Among Caregivers in Four Health Facilities in Adamawa State

Data Collection Tool/Instrument

A validated and pretested semi-structured questionnaire was employed as the data collection tool. Questionnaires were interviewer administered, and baseline data was obtained from all 351 respondents. The questionnaire was a modified version of those used by [17], [18] and [19]. It consisted of three sections: Section 1 on socio-demographic variables, Section 2 on knowledge of Caregivers on Measles and MCV2 and Section 3 on Attitude of Caregivers towards Measles. Reliability test for knowledge gave a Cronbach's alpha of **0.621** and that of attitude gave a Cronbach's alpha of 0.742.

Data Analyses

Data was analysed using Epi Info 7™ and Microsoft Excel, test of normality was conducted to ascertain the distribution of data which will determine the statistical test to conduct; Descriptive analysis was conducted where frequencies, proportions, means and standard deviations of variable were determined. Comparability of the variables was

assessed on the baseline characteristics of the study participants randomized to the intervention and usual care groups, using Chi-square tests. Log-binomial regression was used to estimate the relative risk (RR), difference in risk (RD).

Grading of Caregiver's Knowledge and Attitude

Knowledge

Eight questions were asked on knowledge of Measles and MCV2; each correct answer was graded '1' and wrong answer zero (0); Score range was therefore 0-8.

Attitude

Seven (7) variables were assessed for caregivers' attitude towards MCV2; all questions asked were in the negative perspective so the scores were given as: —Strongly agree; (0 point), —Agreed; (1 point), Disagreed; (2 points) and —Strongly disagreed; (3 points) giving a range of scores of 0-18.

This grading was adapted from previous study [19].

Definition of Terms

Good Knowledge

Caregivers with scores 5-8 (63-100%) are categorized as having good knowledge of measles and MCV2. -rephrase.

Poor Knowledge

A caregiver is categorized as having poor knowledge when he/she scores 0-4 (0-38%).

Positive Attitudes

Caregivers with scores 9-18 (50 -100%) are categorized as having positive attitude towards MCV2 -rephrase.

Negative Attitude

Caregivers with scores 0-8 (0 - 44%) are categorized as having negative attitude towards MCV2.

Ethical Consideration

Approval was sought from the Research Ethical Committee of the Adamawa State Ministry of Health, informed written consent of the participant in the study was obtained, respondents were assured of anonymity and confidentiality of information supplied at all levels of data handling and mothers in control arm were health educated on MCV2 vaccination on completion of data collection even after scheduled time of MCV2 uptake.

Results

Table 1. Sociodemographic Characteristics of Caregivers Surveyed on MCV 2 in Adamawa State

Variables	Frequency; n (%)			Total	Test type	p-value
	Intervention 1	Intervention 2	Control Group			
Age group (years)						
15-24	17 (14.53)	21 (17.95)	17 (14.53)	55 (15.67)	X ²	0.025
25-34	71 (60.68)	58 (49.57)	64 (54.70)	193 (54.99)		
≥35	29 (24.79)	38 (32.48)	36 (30.77)	103 (29.34)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	351 (100.00)		
Mean (SD)	28.83 (2.90)	28.60 (2.81)	29.13 (3.33)	28.85 (3.01)		
95% CI	26.82 – 30.12	21 (17.95)	27.32–31.82	26.24–31.02		
Health Facility						
Gweda Mallam	34 (29.06)	31 (26.50)	34 (29.06)	99 (28.21)	X ²	0.854
Jambutu	26 (22.22)	35 (29.91)	26 (22.22)	87 (24.79)		
Major Aminu	29 (24.79)	25 (21.37)	27 (23.08)	81 (23.08)		
Ngbalang	26 (22.22)	26 (22.22)	30 (25.64)	84 (23.93)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	351 (100.00)		
Relationship with Child						
Aunt	8 (6.84)	16 (13.68)	13 (11.11)	37 (10.58)	F	0.509
Father	1 (0.85)	2 (1.71)	2 (1.71)	5 (1.42)		
Grandmother	0 (0.00)	1 (0.85)	1 (0.85)	2 (0.57)		
Mother	107 (91.45)	98 (83.76)	100 (85.47)	305 (86.89)		
Sister	1 (0.58)	0 (0.00)	1 (0.85)	2 (0.57)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	351 (100.00)		
Caregiver’s educational level						
Informal Edu	9 (7.69)	9 (7.69)	5 (4.27)	23 (6.55)	X2	0.912
Primary	23(19.66)	22 (18.80)	21 (17.95)	66 (18.80)		
Secondary	56 (47.86)	53 (45.30)	57 (48.71)	166 (47.29)		

Tertiary	29 (24.79)	33 (28.21)	34 (29.06)	96 (27.35)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	351 (100.00)		
Religion						
Christianity	54 (46.15)	44 (37.61)	55 (47.01)	153 (43.59)	X ²	0.276
Muslim	63 (53.85)	73 (62.39)	62 (52.99)	198 (56.41)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	351 (100.00)		
Marital Status						
Married	95 (81.20)	98 (83.76)	84 (71.79)	277 (78.92)	F	0.070
Separated/Divorced	0 (0.00)	0 (0.00)	3 (2.56)	3 (0.85)		
Single	22 (18.8)	19 (16.24)	29 (24.79)	70 (19.94)		
Widow	0 (0.00)	0 (0.00)	1 (0.85)	1 (0.28)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	351 (100.00)		
Occupation						
Business	35 (29.91)	31 (26.50)	39 (33.33)	105 (29.91)	F	0.385
Civil Servant	12 (10.26)	16 (13.68)	15 (12.82)	43 (12.25)		
Farmer	5 (4.27)	1 (0.85)	3 (2.56)	9 (2.56)		
Housewife	62 (52.99)	69 (58.97)	60 (51.28)	191 (54.42)		
Student	3 (2.56)	0 (0.00)	0 (0.00)	2 (0.57)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	117 (100.00)		
Place of Child Delivery						
Health Facility	98 (83.76)	105 (89.74)	103 (88.03)	306 (87.18)	F	0.622
Home	18 (15.38)	11 (9.40)	12 (10.26)	41 (11.68)		
Traditional Birth Attendant	1 (0.85)	1 (0.85)	2 (1.71)	4 (1.14)		
Total	117 (100.00)	117 (100.00)	117 (100.00)	117 (100.00)		

A total of 351 caregivers were recruited for the study. Mean age of caregivers across the three study arms was 28.85 (SD±3.01) and ranged 18-60 years. Majority of caregivers were mothers (86.9%) and married (78.9%) across the three study arms; About half were Muslim (56.4), housewives (54.4) and with

have secondary education (47.29). Majority of children brought for MCV1 were born in health facility (87.5%), are female (53.3%), from a family size of greater than 5 persons (64.1%) and living at a distance of 15-30km (70.7%) from health facility (Table 1).

Table 2. Knowledge of Caregivers on Measles and MCV2 in Adamawa State

Knowledge Questions	Correct Response (%)	Incorrect Response (%)
Heard about measles	299 (85.2)	52 (14.8)
Knowledge of at least 3 symptoms of measles	122 (34.8)	229 (65.2)
What causes measles (Measles virus)	55 (15.7)	296 (84.3)
Aware of measles vaccine	63(17.9)	288 (82.1)

Heard and know about the MCV2	123 (35.0)	228 (65.0)
Age at which MCV2 is given (15 months)	211 (60.1)	140 (39.9)
Benefits of MCV2 (Prevent a child from measles infection)	145 (41.3)	206 (58.7)
Knowledge one potential side effects of the MCV2 vaccine	123 (35.1)	228 (64.9)

Significant majority (85.2%) had heard about measles before, while 14.8% had not. Knowledge of the cause of measles revealed that only 15.7% correctly identified the measles virus as the cause, while 28.5% were incorrect, and 55.8% gave partially correct responses. Awareness of the existence of a vaccine to

prevent measles was high, with 82.1% affirming this knowledge, and 17.9% being unaware. Similarly, knowledge about the Measles-Containing Vaccine (MCV2) was substantial, with 65.0% having heard about it, compared to 35.0% who had not (Table 2).

Table 3. Attitude of Caregivers towards Measles and MCV2 in Adamawa State

Attitude Questions	Agree	Disagree	Neutral	Strongly Agree	Strongly Disagree
MCV2 vaccine is important	221(63.0)	3 (0.9)	69 (19.7)	44(12.5)	14 (4.0)
Trust the safety and effectiveness MCV2	218 (62.1)	9 (2.6)	71 (20.2)	42 (12.1)	11 (3.1)
Vaccines have side effect	174(49.6)	32 (9.1)	114(32.5)	9 (2.6)	22 (6.3)
Would recommend MCV2 to other caregivers?	239 (68.1)	3 (0.9)	27 (7.7)	75 (21.4)	7 (2.0)
Every child should receive two doses of measles vaccine	251 (71.5)	0 (0.0)	20 (5.7)	72 (20.5)	8 (2.3)
Measles vaccine can prevent measles outbreak	243 (69.2)	1 (0.3)	17 (4.8)	87 (24.8)	3 (0.9)

A significant majority of respondents (75.5%) believe that the MCV2 vaccine is important, with 63.0% and only a small fraction disagreed (0.9%). 74.1% of respondents expressed confidence in the safety and effectiveness of the MCV2 vaccine with 62.1% agreeing and 2.6% disagreeing. Nearly half (49.6%) of respondents agreed that too many vaccines could have side effects and stress their children while 2.6% strongly disagreed. The

willingness to recommend the MCV2 vaccine to other caregivers was high, with 68.1% agreeing and 21.4% strongly agreeing. The belief that every child should receive two doses of the measles vaccine was widely held, with 71.5% strongly agreeing and 2.3% strongly disagreeing. The belief that the measles vaccine can prevent measles outbreaks in the community was strong, with 69.2% agreeing and only 0.3% disagreed (Table 3).

Table 4. Association Between Sociodemographic Factors and Caregiver's Knowledge on MCV2 in Adamawa State

Variable	Knowledge		Total	χ^2	p-value
	Good	Poor			
Relationship with child	22 (12.0)	15 (10.8)	37 (10.5)	7.67	0.175
Father	5 (2.7)	0 (0.0)	5 (1.4)		
Grandmother	0 (0.0)	2 (1.4)	2 (0.6)		
Mother	183 (86.3)	122 (87.8)	305 (86.9)		
Sister	1 (0.5)	0 (0.0)	1 (0.3)		
Sister in-law	1 (0.5)	0 (0.0)	1 (0.3)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Marital status					
Married	188 (88.7)	89 (64.0)	277 (78.9)	37.0	<0.001
Separated/Divorced	3 (1.4)	0 (0.0)	3 (0.9)		
Single	21 (9.9)	49 (35.3)	70 (19.9)		
Widow	0 (0.0)	1 (0.7)	1 (0.3)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Religion					
Christianity	66 (31.1)	87 (62.6)	153 (43.6)	33.8	<0.001
Muslim	146 (68.9)	52 (37.4)	198 (56.4)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Caregiver educational level					
No Formal Education	15 (7.1)	8 (5.8)	23 (6.6)	6.4	0.094
Primary	34 (16.0)	32 (23.0)	66 (18.8)		
Secondary	96 (45.3)	70 (50.4)	166 (47.3)		
Tertiary	67 (31.6)	29 (20.9)	96 (27.4)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Occupation					
Businesswoman	61 (28.8)	44 (31.7)	105 (29.9)	6.56	0.255
Civil Servant	29 (13.7)	14 (10.1)	43 (12.3)		
Farmer	8 (3.8)	1 (0.7)	9 (2.6)		
Housewife	111 (52.4)	80 (57.6)	191 (54.4)		
Nil	1 (0.5)	0 (0.0)	1 (0.3)		
Student	2 (0.9)	0 (0.0)	2 (0.6)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Residence					
Rural	11 (5.2)	1 (0.7)	12 (3.4)	5.21	0.074
Semi-urban	59 (27.8)	38 (27.3)	97 (27.6)		
Urban	142 (67.0)	100 (71.9)	242 (68.9)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Sex of Child					
Female	115 (54.2)	72 (51.8)	187 (53.3)	0.202	0.653
Male	97 (45.8)	67 (48.2)	164 (46.7)		

Total	212 (100.0)	139 (100.0)	351 (100.0)		
Birth order of Child					
First	21 (9.9)	22 (15.8)	43 (12.3)	4.48	0.214
Second	52 (24.5)	24 (17.3)	76 (21.7)		
Third	68 (32.1)	46 (33.1)	114 (32.5)		
Fourth and above	71 (33.5)	47 (33.8)	118 (33.6)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Place of Child delivery					
Health Facility	181 (85.4)	125 (89.9)	306 (87.1)	2.21	0.331
Home	29 (13.7)	12 (8.6)	41 (11.7)		
Traditional Birth Attendant's	2 (0.9)	2 (1.4)	4 (1.1)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Distance of caregiver house to vaccination site					
15-30 Minutes	159 (75.0)	89 (64.0)	248 (70.7)	17.7	<0.001
Less than 15 Minutes	16 (7.5)	2 (1.4)	18 (5.1)		
More than 30 Minutes	37 (17.5)	48 (34.5)	85 (24.2)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Family size					
Greater than 5	59 (27.8)	67 (48.2)	126 (35.9)	15.1	<0.001
≤ 5	153 (72.2)	72 (51.8)	225 (64.1)		
Total	212 (100.0)	139 (100.0)	351 (100.0)		
Caregivers age group					
15 - 24	41 (19.3)	24 (17.3)	65	27.1	<0.001
25 - 34	130 (61.3)	53 (38.1)	183		
≥ 35	41 (19.3)	62 (44.6)	103		
Total	212 (100.0)	139 (100.0)	351		

Marital status showed a significant association with knowledge level ($\chi^2 = 37$, $p < 0.001$). Among respondents with good knowledge, 88.7% were married, compared to 64.0% with poor knowledge. Religion also exhibited a significant association with knowledge level ($\chi^2 = 33.8$, $p < 0.001$). A greater proportion of respondents with good knowledge were Muslims (68.9%) compared to those with poor knowledge (37.4%). The distance of the caregiver's house to the vaccination site was significantly associated with knowledge level ($\chi^2 = 17.7$, $p < 0.001$). Respondents living 15-30 minutes from the

vaccination site had better knowledge (75.0%) than those living more than 30 minutes away (17.5%). Family size was also significantly associated with knowledge level ($\chi^2 = 15.1$, $p < 0.001$). Among respondents with good knowledge, 72.2% had a family size of 5 or fewer, compared to 51.8% of those with poor knowledge. Caregiver's age group showed a significant relationship with knowledge level ($\chi^2 = 27.1$, $p < 0.001$). The majority of caregivers with good knowledge were in the 25-34 age group (61.3%), whereas a higher proportion of those with poor knowledge were 35 years or older (44.6%). (Table 4).

Table 5. Association Between Sociodemographic Factors and Caregiver's Attitude Towards MCV2 in Adamawa State

Variable	Attitude		Total	χ^2	p-value
	Good	Poor			
Relationship with child					
Aunt	21 (9.0)	16 (13.6)	37 (10.5)	5.36	0.374
Father	5 (2.1)	0 (0.0)	5 (1.4)		
Grandmother	1 (0.43)	1 (0.8)	2 (0.6)		
Mother	204 (87.6)	101 (85.6)	305 (86.9)		
Sister	1 (0.43)	0 (0.0)	1 (0.3)		
Sister in-law	1 (0.43)	0 (0.0)	1(0.3)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Marital status					
Married	207 (88.8)	70 (59.3)	277 (78.9)	49.0	<0.001
Separated/Divorced	3 (1.3)	0 (0.0)	3 (0.9)		
Single	22 (9.4)	48 (40.7)	70 (19.9)		
Widow	1 (0.4)	0 (0.0)	1 (0.3)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Religion					
Christianity	64 (27.5)	89 (75.4)	153 (43.6)	73.3	<0.001
Muslim	169 (72.5)	29 (24.6)	198 (56.4)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Caregiver educational level					
No Formal Education	21 (9.0)	2 (1.7)	23 (6.6)	7.6	0.056
Primary	45 (19.3)	21 (17.8)	66 (18.8)		
Secondary	104 (44.6)	62 (52.5)	166 (47.3)		
Tertiary	63 (27.0)	33 (28.0)	96 (27.4)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Occupation					
Businesswoman	62 (26.6)	43 (36.4)	105 (29.9)	11.1	0.049
Civil Servant	25 (10.7)	18 (15.3)	43 (12.3)		
Farmer	9 (3.9)	0 (0.0)	9 (2.6)		
Housewife	134 (57.5)	57 (48.3)	191 (54.4)		
Nil	1 (0.4)	0 (0.0)	1 (0.3)		
Student	2 (0.9)	0 (0.0)	2 (0.6)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Residence					
Rural	12 (5.2)	0 (0.0)	12 (3.4)	7.2	0.027
Semi-urban	67 (28.8)	30 (25.4)	97 (27.6)		
Urban	154 (66.1)	88 (74.6)	242 (68.9)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Sex of Child					
Female	127 (54.5)	60 (60.8)	187 (53.3)	0.4	0.516
Male	106 (45.5)	58 (49.2)	164 (46.7)		

Total	233 (100.0)	118 (100.0)	351 (100.0)		
Family size					
Greater than 5	64 (27.5)	62 (52.5)	126 (35.9)	21.4	<0.001
≤ 5	169 (72.5)	56 (47.5)	225 (64.1)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Caregivers age group					
15 - 24	49 (21.0)	16 (13.6)	65 (18.5)	36.6	<0.001
25 - 34	140 (60.1)	43 (36.4)	183 (52.1)		
≥ 35	44 (18.9)	59 (50.0)	103 (29.3)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Birth order					
First	30 (12.9)	13 (11.0)	43 (12.3)	8.2	0.043
Second	60 (25.8)	16 (13.6)	76 (21.7)		
Third	72 (30.9)	42 (35.6)	114 (32.5)		
Fourth and above	71 (30.5)	47 (39.8)	118 (33.6)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Place of Child delivery					
Health Facility	196 (84.1)	110 (93.2)	306 (87.2)	8.4	0.04
Home	33 (14.2)	8 (6.8)	41 (11.7)		
Traditional Birth Attendant's	4 (1.7)	0 (0.0)	4 (1.1)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Distance of caregiver house to vaccination site					
15-30 Minutes	175 (75.1)	73 (61.9)	248 (70.7)	17.0	<0.001
Less than 15 Minutes	16 (6.9)	2 (1.7)	18 (5.1)		
More than 30 Minutes	42 (18.0)	43 (36.4)	85 (24.2)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		

Marital status had a significant association with attitude towards MCV2 ($\chi^2 = 49$, $p < 0.001$). Among those with a positive attitude, 88.8% were married compared to 59.3% of those with a poor attitude. A larger proportion of caregivers with a positive attitude were Muslims (72.5%). Caregiver's occupation also showed a significant relationship with attitude ($\chi^2 = 11.1$, $p = 0.049$). Housewives were more likely to have a positive attitude (57.5%) compared to a poor attitude (48.3%). Residence showed a significant association with attitude

($\chi^2 = 7.21$, $p = 0.027$). Family size was significantly associated with attitude towards MCV2 ($\chi^2 = 21.4$, $p < 0.001$). Caregiver's age group showed a significant relationship with attitude ($\chi^2 = 36.6$, $p < 0.001$). Birth order was also significantly associated with attitude ($\chi^2 = 8.17$, $p = 0.043$). The place of child delivery also showed a significant relationship with attitude ($\chi^2 = 6.43$, $p = 0.04$). Distance from the caregiver's house to the vaccination site was significantly associated with attitude ($\chi^2 = 17$, $p < 0.001$). (Table 5).




Figure 2. Correlation Between Knowledge and Attitude of Caregivers Towards MCV2 in Adamawa State

Figure 2 demonstrates a strong and statistically significant positive correlation between the level of knowledge and attitude towards the MCV2 vaccine among caregivers.

This underscores the importance of educational interventions to improve knowledge, which in turn is likely to enhance positive attitudes towards vaccination.

Table 6. Association Between Sociodemographic Factors and Caregiver's Baseline Attitude Towards MCV2 in Adamawa State (2)

Variable	Attitude		Total	χ^2	p-value
	Positive	Negative			
Sex of Child					
Female	127 (54.5)	60 (60.8)	187 (53.3)	0.4	0.516
Male	106 (45.5)	58 (49.2)	164 (46.7)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
family size					
Greater than 5	64 (27.5)	62 (52.5)	126 (35.9)	21.4	<0.001
≤ 5	169 (72.5)	56 (47.5)	225 (64.1)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Caregivers age group					
15 - 24	49 (21.0)	16 (13.6)	65 (18.5)	36.6	<0.001
25 - 34	140 (60.1)	43 (36.4)	183 (52.1)		
≥ 35	44 (18.9)	59 (50.0)	103 (29.3)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Birth order					
First	30 (12.9)	13 (11.0)	43 (12.3)	8.2	0.043
Second	60 (25.8)	16 (13.6)	76 (21.7)		
Third	72 (30.9)	42 (35.6)	114 (32.5)		
Fourth and above	71 (30.5)	47 (39.8)	118 (33.6)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		
Place of Child delivery					
Health Facility	196 (84.1)	110 (93.2)	306 (87.2)	8.4	0.04
Home	33 (14.2)	8 (6.8)	41 (11.7)		
Traditional Birth Attendant's	4 (1.7)	0 (0.0)	4 (1.1)		

Total	233 (100.0)	118 (100.0)	351 (100.0)		
Distance of caregiver house to Health Facility					
15-30 Minutes	175 (75.1)	73 (61.9)	248 (70.7)	17.0	<0.001
Less than 15 Minutes	16 (6.9)	2 (1.7)	18 (5.1)		
More than 30 Minutes	42 (18.0)	43 (36.4)	85 (24.2)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		

The analysis identifies significant association of timely practices with marital status, place of residence, and the age group of caregivers, all of which are associated with levels of knowledge. (Table 6).

Table 7. Association Between Sources of Information on MCV2 and Knowledge of Caregivers on MCV2 in Adamawa State May 2024

Sources of Information about MCV2 vaccine	Attitude		Total	χ^2	p-value
	Good	Poor			
Friends/Family	12 (5.2)	16 (13.6)	28 (8.0)	10.4	0.005
Healthcare Provider	212 (91.0)	93 (78.8)	305 (86.9)		
Media	9 (3.9)	9 (7.6)	18 (5.1)		
Total	233 (100.0)	118 (100.0)	351 (100.0)		

The table examines the relationship between the source of information about the importance of the Measles-Containing Vaccine (MCV2) and caregivers' attitudes. The analysis reveals a statistically significant association between the source of information and attitude towards the vaccine ($\chi^2 = 10.4$, $p = 0.005$). Among caregivers with a good attitude towards the MCV2 vaccine, 91.0% reported learning about the vaccine from healthcare providers, compared to 78.8% of those with a poor attitude. Friends and family were a more common source of information among those with a poor attitude (13.6%) compared to those with a good attitude (5.2%). Information from media sources was cited by 3.9% of those with a good attitude and 7.6% of those with a poor attitude. (Table 7).

Discussion

Measles is an acute and highly contagious viral disease that continues to pose a public health concern, particularly in developing countries. As such, several studies have been conducted to assess the knowledge and attitudes of caregivers towards measles and vaccination

in general. These studies aim to better understand the factors that influence vaccine acceptance and to inform interventions that can increase vaccination coverage.

The study showed that majority of caregivers were mothers who were married. This finding highlights the caring attitude of mothers compared to other categories of caregivers. The study provides valuable insights into the roles and responsibilities of most women play in the lives of their children. This is consistent with other findings in Nigeria [19, 20].

Association between sociodemographic variables such as marital status, religion, distance to vaccination sites, family size, and caregiver's age group showed significant association with the level of knowledge about measles and the MCV2 vaccine among caregivers. This underscores the need for targeted educational interventions that consider these demographic and socio-economic factors to improve vaccination knowledge and coverage. This is in congruent to findings of a study in Abeokuta, Nigeria [18] and a study conducted in Pakistan [21].

Marital status, religion, occupation, residence, family size, caregiver's age group, birth order, place of child delivery, and distance to vaccination site were all significantly associated with caregivers' attitudes towards the MCV2 vaccine which is consistent with findings from study in Edo State Nigeria [22] and in Enugu State Nigeria [23] where these variables were shown to be positively associated with mother's attitude towards measles vaccination; however, this is in variance with findings in France[24] where these factors are not significantly associated with attitude. These findings underscore the importance of considering these demographic and socio-economic factors in designing interventions aimed at improving attitudes towards vaccination.

In addition, area of residence was found to be significantly associated with positive attitude towards MCV2. Caregivers who reside in Urban areas are more likely to have positive attitude towards MCV2 than those who reside in rural areas. This was similar to findings in other studies, where most caregivers that reside in urban areas had positive attitude than other caregiver[17,19]s. The occupation of caregivers was also significantly associated with positive attitude towards MCV2. Caregivers that are housewives were more likely to have a positive attitude towards MCV2 than caregivers with other occupations. This was consistent with findings from other studies, where majority of housewives had a positive attitude towards MCV2 than those of other occupations [18].

Healthcare providers are the primary source of information for caregivers with good knowledge about MCV2. The significant association indicates that caregivers who learn about the vaccine from healthcare providers tend to have better knowledge. This is in consonance with findings from other studies in Nigeria [20] and in Myanmar[25] which revealed that majority of mothers obtained information on the vaccination program

primarily from the health facilities; though this conflicts with findings from Edo State Nigeria [22] where the media was identified as the major source of information on vaccination for mothers. This highlights the crucial role of healthcare providers in disseminating accurate information about MCV2. Efforts to improve knowledge levels should focus on strengthening the role of healthcare providers in public health education. The significant association indicates that caregivers who learn about the vaccine from healthcare providers are more likely to have a positive attitude. This highlights the important role of healthcare providers in shaping positive attitudes towards vaccination. Efforts to improve attitudes should focus on enhancing the involvement of healthcare providers in public health education.

The survey results highlight relatively high level of awareness and partial understanding of measles and the MCV2 vaccine among caregivers in Adamawa State. However, there is a notable proportion of partially correct and incorrect answers, which could be as result of measles being a common disease indicating areas where public health education and awareness campaigns could be improved to enhance comprehensive knowledge about measles and vaccination. The relatively high level of awareness could be as a result the disease' endemicity in the region, however partial understanding of MCV2 could be due to the fact that the second dose was just recently introduced into the expanded program on immunization in Nigeria. This is similar to findings from other studies Nigeria [18, 19].

The study further reveals a high level of positive attitude towards MCV2 among respondents, with strong beliefs in its importance, safety, effectiveness, and role in preventing measles outbreaks. However, concerns about vaccine side effects and the impact of multiple vaccinations on children were also evident, indicating areas where further education and reassurance may be beneficial. This is similar with findings from

other studies which showed positive attitudes of mothers towards measles vaccination [26][23,24]. Nevertheless, some studies carried out in Bauchi and Cross rivers States Nigeria revealed negative attitudes towards vaccination among the mothers [27].

This study also demonstrates a strong and statistically significant positive relationship between the level of knowledge and attitude towards the MCV2 vaccine among caregivers. This underscores the importance of educational interventions to improve knowledge, which in turn is likely to enhance positive attitudes towards vaccination. This corroborates other studies conducted in Nigeria [19, 23]. Furthermore, caregivers with positive attitude towards MCV2 were significantly more likely to vaccinate their children compared to those with a negative attitude which is similar to findings in other studies in Osun State, Nigeria [20] and in Switzerland [28]. This underscores the importance of fostering positive attitudes towards vaccination to improve vaccine uptake and coverage.

Conclusion

This study shows that the knowledge of caregivers on measles and MCV2 is inadequate. There is partial understanding of measles and MCV2 among caregivers in Adamawa State; however, there is relatively high level of awareness on measles with high level of positive attitude towards MCV2 among caregivers.

Major factors associated with good knowledge of measles and MCV2 among caregivers in Adamawa State include age (caregiver being older than 35years), marital status (being married), religion (being a Muslim), marital status (being married) and

distance of caregivers' house to vaccination site (less than 30 minutes' walk to vaccination site). Factors associated with positive attitude of mothers towards MCV2 include marital status (being married), religion (being a Muslim), area of residence (living in urban area), distance of caregivers' house to vaccination site (less than 30 minutes' walk to vaccination site), Birth order (4th order and above) and occupation of caregiver (being a housewife).

A multi-faceted approach is essential to achieve the vaccination coverage needed to eliminate measles in Nigeria. It is essential to continue to monitor and address knowledge gaps and negative attitudes towards measles vaccination to effectively combat measles. Improving vaccination coverage of MCV2 requires increasing awareness of measles and MCV2 through monitoring and addressing knowledge gaps and negative attitudes towards measles vaccinations.

Acknowledgement

The authors extend their sincere gratitude to God for his grace and help throughout the course of this study. Our appreciation goes to the research committee, secretariat members, thesis guidelines formulation committee, and research scholars of Texila American University for their assistance in preparing these guidelines. Special thanks to the Texila University Coordinator for their tireless efforts in bringing this research project to completion. Lastly, we thank all the caregivers for their invaluable contribution to this study.

Declaration of Conflicting Interest

The Author(s) declares that there is no conflict of interest.

References

[1]. Franconeri, L., Antona, D., Cauchemez, S., Lévy-Bruhl, D., Paireau, J., 2019, Two-dose measles vaccine effectiveness remains high over

time: A French observational study 2017–2019, *Vaccine* 2023, 41(39):5797-5804, <https://pubmed.ncbi.nlm.nih.gov/37586955/>

[2]. World Health Organization, 2021, Immunization coverage, Fact Sheet 2021. Retrieved

- From <https://www.who.int/news-room/fact-sheets/detail/immunization-coverage>
- [3]. World Health Organization, 2021, Implementing the Immunization Agenda 2030, WHO 2021:1–32, Retrieved from <https://www.who.int/publications/m/item/implementing-the-immunization-agenda-2030>
- [4]. Mehedi, H. M., Soares, M. R., Ahmed, S., et al, 2021, Meeting the global target in reproductive, maternal, newborn, and child health care services in low- And middle-income countries, *Glob Health Sci Pract.*, 8(4):654-665, <https://pubmed.ncbi.nlm.nih.gov/33361233/>
- [5]. World Health Organization, 2020, Immunization Agenda 2030: a global strategy to leave no one behind, Retrieved from <https://www.who.int/publications/m/item/immunization-agenda-2030-a-global-strategy-to-leave-no-one-behind/> (accessed March 17, 2024)
- [6]. Ward, J. L., Azzopardi, P. S., Francis, K. L., Santelli, J. S., Skirbekk, V., Sawyer, S. M., et al., 2019, Global, regional, and national mortality among young people aged 10–24 years, 1950–2019: a systematic analysis for the Global Burden of Disease Study 2019, *The Lancet* 2021, 398 (10311), 1593 – 1618 [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)01546-4/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)01546-4/fulltext)
- [7]. Nigeria Economic Summit Group, 2021, Health Vaccine Policy, Received from https://nesgroup.org/download_policy_drafts/Nigeria%20Vaccine%20Policy%20%282021%29_Draft%20I_1661875592.pdf
- [8]. World Health Organization, 2021, Development of Vaccines, Retrieved from <https://www.googleadservices.com/pagead/aclk?sa=L&ai=DChcSEwjPIpBwnqaIAxXWl4MHHWBpAPcYABAAGgJlZg&co=1&ase=2&gclid=CjwKC Aj>
- [9]. Wariri, O., Nkereuwem, E., Erundu, N. A., Edem, B., Nkereuwem, O. O., Idoko, O. T., et al., 2021, A scorecard of progress towards measles elimination in 15 west African countries, 2001–19: a retrospective, multi-country analysis of national immunisation coverage and surveillance data, *Lancet Glob Health*, 9(3), e280-e290 <https://pubmed.ncbi.nlm.nih.gov/33607028/>
- [10]. Adamawa State Government, 2020, Adamawa State Situation Analysis of Children, Retrieved from <https://adspc.ad.gov.ng/wp-content/uploads/2024/01/Adamawa-State-Child-Situation-Analysis-SitAn.pdf>
- [11]. Durrheim, D. N., 2020, Measles eradication - retreating is not an option, *Lancet Infect Dis* 2020; 20(6), e138-e14120, <https://pubmed.ncbi.nlm.nih.gov/32197095/>
- [12]. World Health Organization, 2019, More than 140,000 die from measles as cases surge worldwide, Retrieved from <https://www.who.int/news/item/05-12-2019-more-than-140-000-die-from-measles-as-cases-surge-worldwide>
- [13]. Wang, R., Jing, W., Liu, M., Liu, J., 2021, Trends of the Global, Regional, and National Incidence of Measles, Vaccine Coverage, and Risk Factors in 204 Countries From 1990 to 2019, *Front Med (Lausanne)*, 8(2), 205-207, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8810814/>
- [14]. Utazi, C. E., Aheto, J. M., Wigley, A., Tejedor-Garavito, N., Bonnie, A., Nnanatu, C., et al., 2023, Mapping the distribution of zero-dose children to assess the performance of vaccine delivery strategies and their relationships with measles incidence in Nigeria, *Vaccine* 2023, 41(1), 170-181 <https://www.sciencedirect.com/science/article/pii/S0264410X22014207>
- [15]. Ehlman, D. C., Magoola, J., Tanifum, P., Wallace, A. S., Behumbiize, P., Mayanja, R., et al., 2021, Evaluating a mobile phone–delivered text message reminder intervention to reduce infant vaccination dropout in Arua, Uganda: Protocol for a randomized controlled trial, *JMIR Res Protoc*, 10(2), 17262 –17263, <https://pubmed.ncbi.nlm.nih.gov/33625372/>
- [16]. Lwanga, S. K., Lemeshow, S., 1991, Sample Size Determination in Health Studies. *Biometrics* 1991;47 (2), 36-42, https://tbrieder.org/publications/books_english/lemeshow_samplesize.pdf
- [17]. Dalaba, M. A., Ane, J., Bobtoya, H. S., 2023, Factors contributing to low second dose measles-

rubella vaccination coverage among children aged 18 to 59 months in Bolgatanga Municipality of Ghana: a cross-sectional study, *Journal of Global Health Science*, 5(1), 2671-2923, <https://www.e-jghs.org/pdf/10.35500/jghs.2023.5.e11>

[18]. Oluseye, O. M., Jimoh, N. A., Ogunleye, C. A., 2021, Knowledge and Attitude of Mothers towards Measles, Mumps and Rubella MMR Vaccine in Idi- Aba Community Abeokuta, Nigeria, *Open Journal of Medical Research*, 2(2), 136–148. <https://www.openjournalsnigeria.org.ng/pub/ojmr/202006-259.pdf>

[19]. Taiwo, B., Abidemi, L., 2017, Knowledge, Perception, Practice and Sources of information on Immunization among mothers/Caregivers of Children Aged 12-23 months in Kaduna State, Nigeria, 9(1), 168-170, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5462320/>

[20]. Adedire, E. B., Ajumobi, O., Bolu, O., Nguku, P., Ajayi, I., 2021, Maternal knowledge, attitude, and perception about childhood routine immunization program in Atakumosa-west Local Government Area, Osun State, Southwestern Nigeria. *Pan Afr Med J*, 40(1), 8-10, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9474928/>

[21]. Owais, A., Hanif, B., Siddiqui, A. R., Agha, A., Zaidi, A. K., 2011, Does improving maternal knowledge of vaccines impact infant immunization rates? A community-based randomized-controlled trial in Karachi, Pakistan, *BMC Public Health*, 11(1), 238-239, <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-11-239>

[22]. Uwaibi, N. E., Akhimienho, I. K., 2020, Assessment of knowledge and practice of childhood routine immunization among mothers/caregivers attending primary health care centres in Benin City, Nigeria, *Journal of Applied Sciences and Environmental Management*, 24 (10), 1703–1710, [https://www.semanticscholar.org/paper/Assessment-of-knowledge-and-practice-of-childhood-Uwaibi-](https://www.semanticscholar.org/paper/Assessment-of-knowledge-and-practice-of-childhood-Uwaibi-Akhimienho/89b9ffa2659edf5ffe8a83b4f204ab4a5e1f2959)

[Akhimienho/89b9ffa2659edf5ffe8a83b4f204ab4a5e1f2959](https://www.semanticscholar.org/paper/Assessment-of-knowledge-and-practice-of-childhood-Uwaibi-Akhimienho/89b9ffa2659edf5ffe8a83b4f204ab4a5e1f2959)

[23]. Tagbo, B., Uleanya, N., Nwokoye, I., Eze, J., Omotowo, I., 2012, Mothers' knowledge, perception and practice of childhood immunization in Enugu, *Niger J Paediatr*, 39 (3), 314-318, https://www.researchgate.net/publication/272338782_Mothers'_knowledge_perception_and_practice_of_childhood_immunization_in_Enugu

[24]. Pulcini, C., Massin, S., Launay, O., Verger, P., 2014, Knowledge, attitudes, beliefs and practices of general practitioners towards measles and MMR vaccination in southeastern France in 2012, *Clinical Microbiology and Infection*, 20(1), 38–43, <https://pubmed.ncbi.nlm.nih.gov/23517454/>

[25]. Htike, A. Z., Aung, S. M., Win, O. M., 2020, Maternal Knowledge on Routine Childhood Immunization: A Community Based Cross-sectional Study in Rural Area of Mon State, Myanmar, *Asian Journal of Research in Infectious Diseases*, 3(2), 1–9, <https://journalajrid.com/index.php/AJRID/article/view/54>

[26]. Shaheen, A., Afzal, M., Majeed, I., Amir Gilani, S., 2019, Mother's Knowledge and Attitude Towards Childhood Immunization in a Rural Community, *Global Scientific Journals*, https://www.globalscientificjournal.com/researchpaper/Mother_s_knowledge_and_attitude_towards_childhood_immunization_in_a_rural_community.pdf

[27]. Cockcroft, A., Usman, M. U., Nyamucherera, O. F., Emori, H., Duke, B., Umar, N. A., et al., 2014, Why children are not vaccinated against measles: A cross-sectional study in two Nigerian States, *Archives of Public Health*, 72(1), 48-50, <https://archpublichealth.biomedcentral.com/articles/10.1186/2049-3258-72-48>

[28]. Weiss, C., Schröpfer, D., Merten, S., 2016, Parental attitudes towards measles vaccination in the canton of Aargau, Switzerland: A latent class analysis, *BMC Infect Dis*, 16(1), 400 -405, <https://bmcinfectdis.biomedcentral.com/articles/10.1186/s12879-016-1747-0>