





















- longitudinal study,” *Psychogeriatrics*, vol. 21, no. 4, pp. 540–551, 2021, Doi: 10.1111/psyg.12703.
- [5] N. Salari et al., “Prevalence of stress, anxiety, depression among the general population during the Covid-19 pandemic: a systematic review and meta-analysis,” *Global Health*, vol. 16, no. 1, pp. 1–11, 2020.
- [6] L. García-Fernández, V. Romero-Ferreiro, P. D. López-Roldán, S. Padilla, and R. Rodríguez-Jimenez, “Mental Health in Elderly Spanish People in Times of Covid-19 Outbreak,” *Am. J. Geriatr. Psychiatry*, vol. 28, no. 10, pp. 1040–1045, 2020, Doi: 10.1016/j.jagp.2020.06.027.
- [7] S. M. H. Javadi and N. Nateghi, “Covid-19 and Its Psychological Effects on the Elderly Population,” *Disaster Med. Public Health Prep.*, vol. 14, no. 3, pp. e40–e41, 2020, Doi: 10.1017/dmp.2020.245.
- [8] R. B. Hirst et al., “The 5-HTTLPR long allele predicts two-year longitudinal increases in cortisol and declines in verbal memory in older adults,” *Int. J. Geriatr. Psychiatry*, vol. 35, no. 9, pp. 982–988, 2020, Doi: 10.1002/gps.5319.
- [9] C. B. Cunha and S. M. Opal, “Middle East respiratory syndrome (MERS): A new zoonotic viral pneumonia,” *Virulence*, vol. 5, no. 6, pp. 650–654, 2014, Doi: 10.4161/viru.32077.
- [10] T. P. S. Oei, S. Sawang, Y. W. Goh, and F. Mukhtar, “Using the Depression Anxiety Stress Scale 21 (DASS-21) across cultures,” *Int. J. Psychol.*, vol. 48, no. 6, pp. 1018–1029, 2013, Doi: 10.1080/00207594.2012.755535.
- [11] D. S. Hui, E. I. Azhar, Z. A. Memish, and A. Zumla, “Human Coronavirus Infections—severe acute respiratory syndrome (SARS), Middle East Respiratory Syndrome (MERS), and SARS-CoV-2,” *Encycl. Respir. Med.*, no. January, pp. 146–161, 2022, Doi: 10.1016/b978-0-12-801238-3.11634-4.
- [12] L. A. Eaton and S. C. Kalichman, “Social and behavioral health responses to Covid-19: lessons learned from four decades of an HIV pandemic,” *J. Behav. Med.*, vol. 43, no. 3, pp. 341–345, 2020, Doi: 10.1007/s10865-020-00157-y.
- [13] S. Lin, R. Kantor, and E. Clark, *Coronavirus Disease 2019*, vol. 37, no. 4, 2021. Doi: 10.1016/j.cger.2021.05.001.
- [14] M. A.-K. Karataş, “Covid - 19 Pandemisinin Toplum Psikolojisine Etkileri ve Eğitim Yansımaları,” *J. Turkish Stud.*, vol. Volume 15, no. Volume 15 Issue 4, pp. 1–13, 2020, Doi: 10.7827/turkishstudies.44336.
- [15] J. Roy, R. Jain, R. Golamari, R. Vunnam, and N. Sahu, “Covid-19 in the geriatric population,” *Int. J. Geriatr. Psychiatry*, vol. 35, no. 12, pp. 1437–1441, 2020, Doi: 10.1002/gps.5389.
- [16] J. Gustavsson and L. Beckman, “Compliance to recommendations and mental health consequences among elderly in sweden during the initial phase of the Covid-19 pandemic—a cross sectional online survey,” *Int. J. Environ. Res. Public Health*, vol. 17, no. 15, pp. 1–10, 2020, Doi: 10.3390/ijerph17155380.
- [17] K. Lee, G. C. Jeong, and J. Yim, “Consideration of the psychological and mental health of the elderly during Covid-19: A theoretical review,” *Int. J. Environ. Res. Public Health*, vol. 17, no. 21, pp. 1–11, 2020, Doi: 10.3390/ijerph17218098.
- [18] M. Sabe, N. Zhao, and S. Kaiser, “A systematic review and meta-analysis of the prevalence of cocaine-induced psychosis in cocaine users,” *Prog. Neuro-Psychopharmacology Biol. Psychiatry*, vol. 109, no. January 2021, Doi: 10.1016/j.pnpbp.2021.110263.
- [19] A. Barone et al., “The effects of sustained Covid-19 emergency and restrictions on the mental health of subjects with serious mental illness: A prospective study,” *J. Community Psychol.*, no. April, pp. 1–14, 2022, Doi: 10.1002/jcop.22886.
- [20] B. Lábadi et al., “Psychological well-being and coping strategies of elderly people during the Covid-19 pandemic in Hungary,” *Aging Ment. Heal.*, vol. 26, no. 3, pp. 570–577, 2022, Doi: 10.1080/13607863.2021.1902469.