

Effectiveness of Group CBT with Memory Specificity Training and Self-Distancing in Moderately Depressed Adults

Article by Peter John Sabatelli

PhD Clinical Psychology Student, Texila American University

Email: psabatelli@btinternet.com

Abstract

Memory specificity training (MEST) alone can improve mood associated with depression as found in previous studies. The aim was to build upon and expand on previous MEST and self-distancing (SD) studies by testing the effectiveness and feasibility of seven treatment sessions with a three month follow up of group CBT with MEST and SD in moderately depressed adults in two London Boroughs. Method was to recruit 60 participants from advertisements and a screening process that included administration of questionnaires, the autobiographical memory test (AMT) and Beck depression inventory (BDI-II) and interviewed to confirm they met the inclusion and exclusion criteria, and met symptoms for depression as listed in the DSM-5. Consent was obtained and then block randomising done to ensure a balanced group. There were 55 participants who completed the treatment at all time points and their data and scores on the AMT and BDI-II was used when running a one way repeated ANOVA as this was a within group design with repeated measures, using SPSS 22. The results showed improvements occurred at post-treatment and three month follow up for AMS and BDI-II scores that was statistically significant with large effect sizes at post-treatment and three month follow up. Discussion, the null hypothesis was rejected and the alternative hypothesis was accepted. MEST with self-distancing is an effective and feasible adjunct with CBT in improving memory specificity and mood. Limitations included; the study was not fully blind, no independent therapist, low frequency of supervision for checking manual adherence, no SCID, short follow up period and reduced generalisability. Future research could repeat this study using group CBT as a control, participants recruited from out-patient departments, larger sample size, improve blinding before random allocation, using SCID, frequent supervision and use of an independent therapist.

Key words: Cognitive Behavioural Therapy, Memory Specificity Training, Self-Distancing.

Introduction

There is a need to continue the development or refinement of existing evidence based psychological interventions such as CBT for depression due to the relative stagnation in effectiveness of CBT and other psychosocial interventions for depression [1-5]. A promising treatment is memory specificity training (MEST) that was used alone for moderate depression that worked by reducing overgeneral memories (OGM's) and increasing specific memories (AMS), leading to improved mood [9-11]. Also interesting is a potential treatment technique called self- distancing (SD), where induction studies found participants experienced less distress and improved mood.

The first MEST study consisted of four group sessions delivered to moderately depressed Belgium in-patients, aged 32-55 that resulted in increased AMS and depressive symptom improvements. There was no-follow up, no control or comparison group, the sample size of 10 had seven dropping out and all participants were female [9].

In the second MEST study, 23 Afghan adolescents based in Iran with moderate depression, were randomly assigned, 12 to the MEST group and 11 to the control group with a two months follow up. The MEST group retrieved a higher proportion of specific memories and lower levels of depression than did the control group [10]. In this study the benefits of MEST could have been confounded by subsequent improvement in trauma experiences and hence a commensurate reduction in OGM's. The Impacts of events scale was not re-administered at

post-treatment and follow up that would have helped to dismiss or confirm trauma improvements as a possible confounding factor [18]. Trauma factors associated with OGM's will not be a focus of the research here.

The third MEST study recruited and randomly allocated 32 depressed out-patients, aged 27-56 from Amsterdam to undergo originally four sessions of group MEST. Statistical analysis was done on 26 participants, the majority were women who attended between four to five sessions. The results showed a decrease in depressive symptoms corresponding with an increase in AMS [11]. This study had no control group and a relatively small sample size. All the participants whose depressive symptoms were measured at a three-month follow-up had started a psychological or pharmacological treatment at the time of the follow-up. This makes it hard to distinguish whether further improvement of symptoms measured at follow-up was caused by MEST only.

The SD induction studies was utilised primarily in young non-clinically depressed participants [12-17]. Only one study induced SD on a depressed group with no extended follow up period nor used with any other psychological treatment [12]. In the latter study, they recruited 51 moderately depressed adults and 45 healthy adults without any history of depression in America. They then used cues to recall a past experience in which they felt overwhelming sadness and were randomly assigned to either adopt a self-immersed or a self-distanced perspective to neutral or depressotypic words and recorded the speed of responses. The depressed group that analysed their feelings from a self-distanced perspective showed lower levels of depressive thinking and reduced low mood than the self-immersed group. They suggested that future research on SD should examine whether these findings generalize to adults with depression under other conditions, which could be taken to also apply to treatment conditions[16].

A brief outline of the literature will now be given about identifying depression, impact OGM's has on depression, defining OGM's and AMS, the theoretical framework to explain how OGM's and AMS occur, distancing connection to reframing and AMS, the purpose for the research, significance and hypotheses.

Depression consists of low mood, loss of interest or reduced satisfaction, enjoyment in activities for at least a two week period, along with other symptoms causing distress and diminished functioning [31]. Overgeneral autobiographical memories (OGM) which is considered an avoidance strategy [6-7] are strongly associated with making depression more severe, delays recovery and is a vulnerability factor for subsequent episodes of depression [8].

Overgeneral autobiographical memories (OGM) are modifiable [8] and defined as including a summary of several similar events (e.g., "watching my favourite TV show on Saturday evenings") or a memory of a general time period that spans more than one day (e.g., "My holiday in France last year" [32, 7]. Autobiographical memory specificity (AMS) is defined as a memory for a specific occasion, event that occurred at least one week or longer at a particular place and time within a 24 hour period [9-10] thus relates to spatial-temporal specificity not the content of memory [33].

The main theoretical framework to understand OGM is based on the self-memory system model [20]. It suggests that knowledge is arranged hierarchically along lifetime stages that can be accessed through memories that range from the abstract to the concrete or general events to specific events [8, 20]. It is not the content of thought of the memory that matters but the degree of spatial-temporal specificity and arousing details that is recalled [21]. OGM's lacks spatial-temporal specificity and emotional arousal and AMS has more sensory-perceptual arousing details.

Another more recent model to explain the mechanisms that lead to OGM in depression is from the capture and rumination, functional avoidance, executive control (CaRFAX) model [8, 22]. This model hypothesize that difficulties accessing AMS result from the capture of memory search efforts by the bringing together of OGM's that causes or keeps the depression in place. This is made worse by avoidance of specific aspects of distressing autobiographical events that can become habitual, leading to processing of broader life experiences only at the level of OGM's [8] that can cause of keep the depression in place.

A distancing stance is required to enable reframing in CBT [20, 25] and distancing is an important component in mindfulness for depression [21], both CBT and mindfulness are linked to increasing AMS [22-24]. It is reasonable to assume the distancing element in SD may also help to enhance AMS and may help with facilitating specific re-appraisal as is necessary in CBT [25].

The distancing component of SD, means taking a mental step back. This stance permits a broader context to be observed at a distance when recalling or retrieving past experiences which facilitates the second component of SD, reconstruing. The distancing component may permit flexibility and modification to occur to allow new explanation, understanding, meaning, insights that could reduce current and future distress [13-16].

The research studies above suggested that reducing OGM's and increasing AMS could improve mood and MEST was developed to reduce OGM's and increase AMS with promising results [10-12]. The limitations in MEST studies suggest a larger sample size is required, from a different country, with block randomisation to better control gender allocation and group balance. In addition two relatively recent studies recommended that future studies could use MEST as an adjunct with CBT [14, p.8; 11, p.7]. Another gap in research is that SD could be used with depressed adults under others conditions. Here it could be under treatment conditions such as with CBT with MEST [16].

The purpose of the study was to build and expand on the previous research on MEST and SD. This was done by examining the effectiveness and feasibility of group CBT with MEST and SD using AMS and low mood measures in moderately depressed adults from London, UK, under pre-treatment, post-treatment and three month follow up conditions.

Hypothesis 1: There would be a difference in the total score for AMS by moderately depressed adults by post- treatment and three month follow up conditions as a result of group CBT with MEST and SD. Hypothesis 2: There would be a difference in the total score on the BDI-II by moderately depressed adults by post- treatment and three month follow up conditions as a result of group CBT with MEST and SD.

It is important to seek to improve outcomes [1-5] for moderately depressed adults that could further help to reduce social and economic burdens [28-30]. It could reduce suffering quicker for those who are depressed but also for their families as well and personally reduce health, social and economic care costs. This may be achieved by treating more depressed clients through groups than individual one to one treatment as in group CBT with MEST and SD with specific clinical health benefits of increasing AMS to reduce vulnerability to recurrence of depression [7-8] and improved mood [9-11].

Method

Participants

Participants were 30 male and 30 female adults, aged 24-56 ($M = 36.92$; $SD = 9.28$), recruited from two London Boroughs fulfilled the inclusion and exclusion criteria and consented to treatment. The inclusion criteria stated that participants had to be aged between 20-60, presence of moderate symptoms defined as a Beck depression inventory (BDI-II) score of 20 or more but less than 29 [38] and AMS of less than 0.70 as assessed on the AMT [42]. The exclusion criteria included high levels of suicidality or harm to others as taken from BDI-II scores; secondary diagnosis of another affective disorder or a psychotic disorder; current drug/ alcohol abuse or dependence, personality disorder (assessed via participant report); presence of head trauma or organic brain damage (assessed via participant report); history of childhood abuse; PTSD symptoms, chronic pain, adjustment problem, primary problem is anxiety, poor grasp of the English language, reading and writing.

All were experiencing moderate depression as confirmed by the researcher and other experienced clinician using the DSM-5 major depression criteria [31] and scores based on the BDI-II [38]. The mean score on the BDI-II ($M = 26.23$; $SD = 1.15$) suggested the group had moderate depression.

Data from Raes et al [9] suggested a sample size of 22 (11 per group) would provide 80% power, with a directional alpha of .05, to detect a similar improvement in AMS. The sample

size had to be larger than 11 per group as a requirement of 36 or more for each treatment group was needed to achieve medium effect sizes averaging $d = 0.51$ in previous research on self-distancing [12-13].

Procedure

A manual was compiled for the study, the CBT components consisted of behavioural activation and cognitive reframing that adhered to the Beckian cognitive model and Lewholm behavioural model [25, 39-40]. Both have validity and reliability when used to treat depression [5]. MEST training has some degree of validity, reliability [9-11]. A similar format was used but spread over seven sessions instead of five due to having to accommodate the CBT components and the self-distancing technique. The self-distancing technique that was used has some degree of validity [12-17]. The manual was also checked by several highly experienced clinical psychologists who confirmed the manual developed contained the core components of CBT and broadly the MEST protocol whereas a SD protocol they could only confer face validity.

Ethical approval was obtained then the research was advertised in two London boroughs. Prospective participants responded by e-mail to leaflets posted through the letterbox of people's homes, flyers posted on the advertisement boards in supermarkets, health clubs, local libraries, community centres, religious places of worship, several newsagents and charity shops. Those who responded to the advert were invited to a screening session to assess for treatment suitability.

Participants were then screened using the autobiographical memory test (AMT) and Beck depression Inventory (BDI-II). They were then interviewed to ensure they met the inclusion criteria and had symptoms of depression as listed in the DSM-5 [31]. Consent was obtained when it was satisfied they fully understood the contents of the participants information sheet and understood they could withdraw at any time without prejudice. They were then cluster randomised using a stratified randomisation procedure [41] to ensure a balanced gender group. Those not selected were sent a letter thanking them for attending and wishing them well. A total of 60 participants were recruited this way, five did not start treatment, with 55 completers.

A within subject design was selected. The within group factors, included pre-treatment, post-treatment and three month follow up to examine the effectiveness of group CBT with MEST and SD.

To help guarantee the integrity of each treatment, that the group manual was adhered to, a two random audio recordings and two random in observations were made between session two to seven by an independent clinical psychologist. This was to check for manual adherence and bias in non-verbal presentation, none was found.

Measures

The Autobiographical Memory Test measures both OGM and AMS [42] with a reliability estimate of 0.72 [53]. For example, ten cue-words (five positive and five negative), are provided in turn and participants then write down one AMS that the cue-word reminds them of within 30 or 60seconds. Instructions make it clear AMS refers to one particular occasion or event that happened on a particular day, within 24 hour period and at least one week before the test. Each response is coded as an AMS, a memory for an event, occasion that happened within a 24 hour period on a particular day, more than one week ago. OGM includes extended and categoric memory, were a repeated event or occasion is recalled without specifying any particular time or lasted longer than a 24 hour period or more than one day. The other is considered a verbal association to the cue rather than a memory and coded semantic and lastly no memory or response is given and coded an omission.

In this research, two parallel sets of 18 cue-words, nine positive and nine negative, similar to a previous MEST study [10] were matched for familiarity and emotionality using three independent raters [43] for pre-test, post and follow up.

The BDI-II is a 21-item self-report questionnaire used for measuring the severity of depression in adolescents and adults aged 13 and older. Each response relates to the recent

two week period and for each question a score of 3, 2, 1, or 0 may be selected with total scores ranging from 0 to 63. The questions had been revised to correspond more with the DSM-IV criteria for depression. Total scores ranging from 0 to 13 represent normal to minimal depression; 14 to 19 is considered mild depression; 20 to 28 is seen as moderate and between 29 to 63 as severe or major. It has high reliability and capacity to discriminate between depressed and non-depressed subjects, concurrent, content, and structural validity and internal consistency gave a cronbach's alpha of 0.92 for outpatients and 0.93 for students [39, 44].

Statistical Analysis

In order to determine changes in memory specificity and low mood from pre-treatment to post-treatment and three month follow up, using SPSS 22, a one way repeated ANOVA with a pairwise comparison was run. The statistical significance level for the two hypotheses was set at .01 with a confidence level of 99% for all statistical tests. Effect sizes are indicated in terms of both partial η^2 and reported as significant at $p < .001$.

Results

Sample

Sixty participants were included in the group CBT with MEST intervention that received pre-treatment measures. Five of the participants did not start the intervention due to work commitment, starting treatment elsewhere, illness and carers responsibilities.

Data on the 55 completers were used for data analysis, this equates to 8.3% of missing data in subsequent measures at post-treatment and three month follow up. Missing data was dealt with using the listwise default from the SPSS 22 [45] as the percentage of 8.3% of missing data does not compromise statistical analysis [46-47].

The skewness and kurtosis scores were within plus or minus 2.58 at pre-treatment, post-treatment and three month follow up.; there was no outliers as no residuals $\geq \pm 3$ were obtained at pre-treatment and normal Q-Q plot of residuals, showed they are not too distorted from the diagonal line to suggest that the data did not violate the assumption of normality.

The mauchly's test of sphericity indicated that the assumption of sphericity had been violated for AMS, (2) = 39.102, $p = .001$; and BDI-II, (2) = 17.360, $p = .001$. Epsilon was 0.667 as calculated according to Greenhouse & Geisser [48], and was used to correct the one-way repeated measures ANOVA. The group intervention elicited statistically significant changes in AMS scores over time, $F(1.314, 70.968) = 480.490$, $p < .001$, partial = .989.

Table 1. Baseline characteristics of group sample (n=60) at pre-treatment

Demographic Variables	Category	Group N = %	Mean and SD
Gender	Male	30 = 50%	
	Female	30 = 50%	
Marital Status	Married	23 = 38.3%	
	Single	15 = 25%	
	Separated	12 = 20%	
	Divorced	10 = 16.7%	
6 Ethnicity	White	30 = 50%	
	Asian	14 = 23.3%	
	Caribbean	9 = 15%	
	African	4 = 6.7%	
	Other	3 = 5%	

Religion	C of E	22 = 36.7%	
	Catholic	19 = 31.37%	
	Muslim	11 = 18.3%	
	Hindu	4 = 6.7%	
	Other	4 = 6.7%	
Qualification	One or more	55 = 91.7%	
	None	5 = 8.3%	
Occupation	Employed	31 = 51.7%	
	Unemployed	29 = 48.3%	
Age		24-56 (M = 36.92, SD=9.28)	
AMS			(M=56.65, SD=3.03)
BDI-II			(M=26.23, SD=1.15)

Data are mean \pm standard deviation, unless otherwise stated. There was an increase in AMS from 56.6 ± 3.0 at pre-intervention to 91.3 ± 2.3 at post-intervention, a statistically significant increase of 34.6 (99% CL, 33.0 to 36.2), $p < .001$. Data are mean \pm standard deviation, unless otherwise stated.

There was an increase in AMS from 56.0 ± 2.6 at pre-intervention to 96.3 ± 2.3 at three month follow up, a statistically significant increase of 39.7 (99% CL, 38.1 to 41.3), $p < .001$. Epsilon (ϵ) was 0.782, as calculated according to Greenhouse & Geisser [48], and was used to correct the one-way repeated measures ANOVA. The group intervention elicited statistically significant changes in BDI-II scores over time, $F(1.563, 84.420) = 4536.820$, $p < .001$, partial eta squared = .988.

Data are mean \pm standard deviation, unless otherwise stated. There was a decrease in BDI-II scores from 26.3 ± 1.2 at pre-intervention to 7.9 ± 1.3 at post-intervention, a statistically significant decrease of 18.3 (99% CL, 17.5 to 19.1), $p < .001$. Data are mean \pm standard deviation, unless otherwise stated. There was a decrease in BDI-II scores from 26.3 ± 1.2 at pre-intervention to 3.8 ± 1.7 at three month follow up a statistically significant decrease of 22.5 (99% CL, 21.6 to 23.5), $p < .001$.

Discussion

The statistical results support the effectiveness and feasibility of CBT with MEST and SD in treating moderately depressed adults. It also further confirms that OGMs are modifiable [8] and has built on and expanded the research on MEST [10-12, 11, p.7; 14, p.8] and SD [12]. It is also clinically significant as the scores on the BDI-II indicated the participants were in remission at post-treatment and in recovery at three month follow up [39].

Hypothesis one predicted that there would be a difference in effect between pre-treatment, post-treatment and at three month follow up in AMS scores as a result of group CBT with MEST and SD for moderately depressed adults. The results showed there was statistically significant difference at post-treatment and three month follow up. This result rejected the null hypothesis and supported the alternative hypothesis.

The addition of MEST and SD to CBT appeared to enhance AMS as the scores in this study showed higher rates of improvement compared to previous studies using MEST alone [10-12]. AMS improvement in the first MEST study by Raes et al [9] increased by 0.28 by the end of treatment. In the second MEST study by Neshat et al it was by 0.32 at two months follow up [10] and in the third MEST study there was an improvement in AMS of 0.18 in Eigenhuis et al study[11]. The CBT with MEST and SD had by post-treatment showed that

AMS had improved by 34.5 and at three month follow by 39.9 compared to pre-treatment. The result also challenges the research that asserted that taking an analytical focus as found in SD may not reduce OGM [49] found in depression.

Hypothesis two predicted that there would be a difference in effect between pre-treatment, post-treatment and at three month follow up in BDI-II scores as a result of group CBT with MEST and SD for moderately depressed adults. The results showed there was statistically significant difference at post-treatment and three month follow up. This result rejected the null hypothesis and supported the alternative hypothesis.

There is support that the MEST element did contribute to improved mood through increasing AMS, supporting previous research that used MEST alone for adult moderate depression that it can improve mood [9-11]. This may also be due to SD helping to make it easier to re-appraisal and interpret one's experience or 'reconstrue' [13, 26] helping to increase AMS scores.

Limitations include the way of presenting cues and/or available amount of time to respond to the AMT cue words are moderators of AMT performance [50]. Study was not fully blind; no independent therapists used to run groups, no SCID used; participants were highly motivated as evidenced by voluntary participation, high attendance rate and high homework compliance factors not typical of clinically depressed adults that attend out-patient clinics for CBT [51]. There is reduced generalisability due to participants being solely depressed, not on anti-depressant medication and no co-morbidity. The researcher running the group intervention did not receive training to use the treatment manual, did not receive weekly supervision to check manual adherence [52] when running the groups. This was limited to a few random audio recordings and direct observations by an independent experienced clinician.

Conclusion

The findings of this study expanded the work of previous researchers in the area of MEST for depression [9-11] and SD [12-17, 19]. These results go some way towards showing that CBT with MEST and SD is not only feasible but is effective, 'scalable', cost effective due to being a group treatment, potentially may reduce one risk factor OGM's that is associated depression relapse and potentially allows others to get back to work quicker, lowering personal and government health costs.

Future research can repeat this study by having a larger sample size; blind, using SCID, regular independent supervision for the deliverer of group intervention; using independent therapist to run the group intervention. Samples of participants recruited into a randomised controlled trial that is seen at a typical out-patient clinic in comparing five sessions of MEST with five sessions of group CBT with MEST and SD and five sessions of standard group CBT as a control group to see any enhancing effect on the rate and amount of AMS and mood changes in moderately clinical depressed adult participants.

References

- [1] J. Barth, T. Munder, H. Gerger, E. Nüesch, S. Trelle, H. Znoj, P. Juni, P. Cuijpers, Comparative Efficacy of Seven Psychotherapeutic Interventions for Patients with Depression: A Network Meta-Analysis. *PLoS Medicine*, (2013), 10(5), e1001454. doi:10.1371/journal.pmed.1001454
- [2] P. Cuijpers M. Berking, G. Andersson, L. Quigley, A. Kleiboer A, K.S. Dobson, A meta-analysis of cognitive behavior therapy for adult depression, alone and in comparison to other treatments. *Canadian Journal of Psychiatry*, (2013), 58, 376-385
- [3] E. Driessen, H.L. Van, F.J. Don, J. Peen, S. Kool, D. Westra, M. Hendrikson, M. Schoevers, P. Cuijpers, J. Twisk, J.J. Dekker, The efficacy of cognitive behavioral therapy and psychodynamic therapy in the outpatient treatment of major depression: a randomized clinical trial. *American Journal of Psychiatry* (2013), 170:1041-1050. doi: 10.1176/appi.ajp.2013.12070899
- [4] V. Hunot, T.H.M. Moore, D.M. Caldwell, T.A. Furukawa, P. Davies, H. Jones, M. Honyashiki, P. Chen, G. Lewis, R. Churchill R, 'Third wave' cognitive and behavioural therapies versus other psychological therapies for depression. *Cochrane Database of Systematic Reviews* (2013), Art. No.: CD008704. doi:10.1002/14651858.CD008704.pub2.
- [5] T.J. Johnsen and O. Friborg, The Effects of Cognitive Behavioral Therapy As an Anti-Depressive Treatment is Falling: A Meta-Analysis. *Psychological Bulletin*. (2015), Advance online publication. doi.org/10.1037/bul000015

- [6] F. Raes, D. Hermans, A. De Decker, P. Eelen, J.M.G. Williams, Autobiographical Memory Specificity and affect regulation : an experimental approach. *Emotion* (2003), 3, 201-206, doi:10.1037/1528-3542.3.2.201
- [7] J.M.G. Williams, T. Barnhofer, C. Crane, D. Herman, F. Raes, E. Watkins, T. Dalgleish, Autobiographical memory specificity and emotional disorder. *Psychological Bulletin*, (2007), 133, 122-148. doi: 10.1037/0033-2909.133.1.122
- [8] J.A. Sumner, J.W. Griffith, S. Mineka, S. Overgeneral autobiographical memory as a predictor of the course of depression: A meta-analysis. *Behaviour Research Therapy* (2010), 48, 614-625. doi: 10.1016/j.brat.2010.03.013
- [9] F. Raes, J.M.G. Williams, D. Hermans, Reducing cognitive vulnerability to depression: A preliminary investigation of Memory Specificity Training (MEST) in inpatients with depressive symptomatology. *Journal Behavior Therapy Experimental Psychiatry* (2009), 40, 24-38. doi:10.1016/j.jbtep.2008.03.001
- [10] H.T. NeshatDoost, T. Dalgleish, W. Yule, M. Kalantari, S.J. Ahmadi, A. Dryregrov, L. Jobson, L. Enhancing autobiographical memory specificity through cognitive training: an intervention for depression translated from basic science. *Clinical Psychological Science* (2013), 1:84-92. doi: 10.1177/2167702612454613
- [11] E. Eigenhuis, A. Seldenrijk, A. van Schaik, F. Raes, P. vanOppen, Feasibility and effectiveness of memory specificity training in depressed outpatients: a pilot study. *Clinical Psychology and Psychotherapy*. (2015), doi: 10.1002/cpp.1995
- [12] E. Kross, Ö. Ayduk, W. Mischel, When asking 'why' does not hurt: Distinguishing rumination from reflective processing of negative emotions. *Psychological Science* (2005), 16(9), 709-715. doi:10.1111/j.1467-9280.2005.01600.x
- [13] E. Kross and Ö. Ayduk, Facilitating adaptive emotional analysis :Distinguishing distanced-analysis of depressive experiences from immersed-analysis and distraction. *Personality and Social Psychology Bulletin*, (2008), 34, 924 -938. doi:10.1177/0146167208315938
- [14] E. Kross and Ö. Ayduk, Boundary conditions and buffering effects: Does depressive symptomatology moderate the effectiveness of distanced analysis for facilitating adaptive self-reflection? *Journal of Research in Personality* (2009), 43, 923-927. doi:10.1016/j.jrp.2009.04.004
- [15] Kross, E., and Ayduk, Ö. (2011). Making meaning out of negative experiences by self-distancing. *Current Directions in Psychological Science*, 20(3), 187-191. doi: 10.1177/0963721411408883.
- [16] E. Kross, D. Gard, P. Deldin, J. Clifton, Ö. Ayduk, "Asking Why" From a Distance: Its Cognitive and Emotional Consequences for People With Major Depressive Disorder. *Journal of Abnormal Psychology* (2012), 3, 559 -569. doi: 10.1037/a0028808
- [17] E. Kross, E., Bruehlman-Senecal, J. Park, A. Burson, A. Dougherty, H. Shablack, R. Bremmer, J. Moser, Ö. Ayduk, Self-talk as a regulatory mechanism: How you do it matters. *Journal of Personality and Social Psychology* (2014), 106(2), 304. doi: 10.1037/a0035173
- [18] S.A. Moore and L.A. Zoellner, Overgeneral Autobiographical Memory and Traumatic Events: An Evaluative Review. *Psychological Bulletin* (2007), 419-437. doi: 10.1037/0033-2909.133.3.419
- [19] Ö. Ayduk and E. Kross, From a distance: Implications of spontaneous self- distancing for adaptive self-reflection. *Journal of Personality and Social Psychology* (2010), 98(5), 809-829. doi: 10.1037/a0019205
- [20] A.T. Beck, Cognitive therapy of depression: New perspectives. In P.J. Clayton & J. E. Barrett (Eds.), *Treatment of depression: Old controversies and new approaches*. New York: Raven Press (1983), pp. 265-284
- [21] J. Gecht, R. Kessel, T. Forkmann, S. Gauggel, B. Drueke, A. Scherer, V. Mainz, A mediation model of mindfulness and decentering: sequential psychological constructs or one and the same? *BMC Psychology* (2014), 2:18. doi:10.1186/2050-7283-2-18
- [22] C. McBride, Z. Segal, S. Kennedy, M. Gemar, Changes in autobiographical memory specificity following cognitive behavior therapy and pharmacotherapy for major depression. *Psychopathology* (2007), 40, 147-152
- [23] Z. Ranjbarokhan, H.T. Neshatdoost, H. Molavi, H. M. Maeroofi, A comparison of autobiographical memory specificity changes following cognitive behavioral therapy and memory specificity training (MEST) in patients with major depression. *Journal of Contemporary Research in Business* (2012), 4(3):450-455.
- [24] J.M.G. Williams, J. D Teasdale, Z.V. Segal, J. Soulsby, Mindfulness-based cognitive therapy reduces overgeneral autobiographical memory in formerly depressed patients. *Journal of Abnormal Psychology* (2000), 109, 150-155. doi: 10.1037//0021-843X.109.1.150
- [25] A.T. Beck, J.A. Rush, B.F. Shaw, G. Emery, *Cognitive therapy for depression*. New York: Guilford Press (1979).
- [26] A. Bernstein, Y. Hadash, Y. Lichtash, G. Tanay, K. Shepherd, D.M. Fresco, Decentering and Related Constructs A Critical Review and Metacognitive Processes Model. *Perspective Psychological Science* (2015), 10(5):599-617
- [27] T. Dalgleish, A. Bevan, A. McKinnon, L. Breakwell, V. Mueller, I. Chadwick, S. Schweizer, C. Hitchcock, P. Watson, F. Raes, L., Jobson, L., A. Werner-Seider., A., A comparison of Memory Specificity Training (MEST) to education and support (ES) in the treatment of recurrent depression: study protocol for a cluster randomised controlled trial. *Trials*.(2014), 15, 1-9. doi:10.1186/1745-6215-15-293
- [28] P. McCrone, S. Dhanasiri, A. Patel, M. Knapp, S. Lawton-Smith, Paying the price: the cost of mental health care in England to 2026. *The King's Fund*, London (2008)
- [29] A.J. Ferrari, F.J. Charlson, R.E. Norman, S.B. Patten, G. Freedman, C.L.J. Murray, T. Vos, H.A. Whiteford, Burden of Depressive Disorders by Country, Sex, Age, and Year: Findings from the Global Burden of Disease Study 2010. *PLoS Med* (2013), 10(11): e10001547. doi:101371
- [30] N. Sartorius. The economic and social burden of depression. *Journal Clinical Psychiatry* (2001), 62(15), 8-11.
- [31] American Psychiatric Association. *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing (2013)

- [32] Conway, M. A., and Rubin, D. C. (1993). The structure of autobiographical memory. In A. E. Collins, S. E. Gathercole, M. A. Conway, & P. E. M. Morris (Eds.), *Theories of memory* (pp. 103-137). Hove, Sussex, England: Erlbaum
- [33] Peeters, F., Wessel, I., Merckelbach, H., and Boon-Vermeeren, M. (2002). Autobiographical memory specificity and the course of major depressive disorder. *Comprehensive Psychiatry*, 43, 344-350. doi:10.1053/comp.2002.34635
- [34] M.A. Conway and C.W. Pleydell-Pearce, The construction of autobiographical memories in the self-memory system. *Psychological Review*, (2000), 107, 261–88. doi: 10.1037//0033-295X.107.2.261
- [35] K.M. Davis, M.A. Lau, D.R. Cairns, Development and preliminary validation of a trait version of the Toronto mindfulness scale. *Journal of Cognitive Psychotherapy*, (2009), 23, 185-197. doi:10.1891/0889-8391.23.3.185
- [36] D. M. Fresco, M.T. Moore, M.H.M. van Dulmen, Z.V. Segal, S.H. Ma, J.D., Teasdale, J.M.G. Williams, Initial Psychometric Properties of the Experiences Questionnaire: Validation of a Self-report Measure of Decentering Behavior Therapy (2007), 38, 234-246, <http://dx.doi.org/10.1016/j.beth.2006.08.003>
- [37] L.M. McCracken, E. Barker, E., J. Chilcot, Decentering, rumination, cognitive defusion, and psychological flexibility in people with chronic pain. *Journal Behavioral Medicine* (2014), 37, 1215-1225. doi: 10.1007/s10865-014-9570
- [38] A.T. Beck, R.A. Steer, G.K. Brown, *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation, (1996)
- [39] D. Barlow, *Clinical Handbook of Psychological Disorders, Fifth Edition: A Step-by-Step Treatment Manual*. The Guildford Press, London (2008)
- [40] K. Efthimiou and M. Psoma, Lewinsohn's cognitive behavioral group Therapy course for depression: structure, application and efficacy. *Encephalos*, (2012), 49, 60-66.
- [41] K.P. Suresh, An overview of randomization techniques: An unbiased assessment of outcome in clinical research. *Human Reproductive Science* (2011), 4, 8–11. doi: 10.4103/0974-1208.82352
- [42] J.M.G. Williams and K. Broadbent, Autobiographical memory in suicide attempters. *Journal of Abnormal Psychology* (1986), 95, 144-149. doi:10.1037/0021-843X.95.2.144
- [43] J.W. Griffith, J.A. Sumner, E. Debeer, F. Raes, D. Hermans, S. Mineka, R.E. Zinbarg, M.G. Craske, An item response theory/confirmatory factor analysis of the Autobiographical Memory Test. *Memory* (2009), 17, 609–623.
- [44] Y.P. Wang and C. Gorenstein, Psychometric properties of the Beck Depression Inventory-II: a comprehensive review. *Revista Brasileira de Psiquiatria* (2013), 35:416–431. doi:10.1590/1516-4446-2012-1048
- [45] G.L. Schlomer, S. Bauman, S. N.A. Card, Best Practices for Missing Data Management in Counselling Psychology (2010), Vol. 57, No. 1, 1–10. DOI: 10.1037/a0018082
- [46] D.A. Bennett, How can I deal with missing data in my study? *Australian and New Zealand Journal of Public Health*, (2001), 25, 464 – 469
- [47] C.Y.J. Peng, M. Harwell, S.M. Liou, L.H. Ehman, Advances in missing data methods and implications for educational research. In S. Sawilowsky (Ed.), *Real data analysis* (pp. 31–78). Greenwich, CT: Information Age, (2006)
- [48] S.W. Greenhouse and S. Geisser, On methods in the analysis of profile data. *Psychometrika* (1959), 24, 95–112. doi: 10.1007/BF02289823.
- [49] E. Watkins and J.D. Teasdale, Rumination and overgeneral memory in depression: effects of self-focus and analytic thinking. *Journal of Abnormal Psychology* (2001), 110, 353–357.
- [50] M.F. van Vreeswijk and de E.J. Wilde, de, Autobiographical memory specificity, psychopathology, depressed mood and the use of the AMT: A meta-analysis. *Behaviour Research and Therapy* (2004), 42(6): 731-43
- [51] M. Seligman, The effectiveness of psychotherapy. *American Psychologist* (1995), 50, 965–974.
- [52] P.C. Kendall and R.S. Beidas, *Professional Psychology: Research and Practice* (2007), 38(1), 13-20. <http://dx.doi.org/10.1037/0735-7028.38.1.13>
- [53] The factor structure of the Autobiographical Memory Test in recent trauma survivors. J.W. Griffith, B. Kleim, J.A. Sumner, Jennifer and A. Ehlers (2012), The factor structure of the Autobiographical Memory Test in recent trauma survivors. *Psychological Assessment*, 24(3), 640-646. <http://dx.doi.org/10.1037/a0026510>

Funding

This study was self-funded by the author.

Conflict of Interest

There are no conflicts of interest for the author.