





















in South Africa: a cross-sectional study. *BMC Cancer* 2009, 9(1):275.

[49] Blossom DB, Beigi RH, Farrell JJ, Mackay W, Qadadri B, Brown DR, Rwambuya S, Walker CJ, Kambugu FS, Abdul-Karim FW: Human papillomavirus genotypes associated with cervical cytologic abnormalities and HIV infection in Ugandan women. *J Med Virol* 2007, 79(6):758–765.

[50] Parham GP, Sahasrabudde VV, Mwanahamuntu MH, Shepherd BE, Hicks ML, Stringer EM, Vermund SH (2006): Prevalence and predictors of squamous intraepithelial lesions of the cervix in HIV-infected women in Lusaka, Zambia. *Gynecol Oncol* 2006, 103(3):1017–1022.

[51] Akubue AU, Jalal-Eddeen AS, Abdullahi S (2018). An Evaluation of the Integrated Disease Surveillance and Response (IDSR) in Enugu State, Nigeria. *Journal of Health, Medicine, and Nursing* www.iiste.org ISSN 2422-8419. Vol.48, 2018.

[52] Morhason-Bello Imran O., Kareem Yusuf Olushola, Adewole Isaac F. (2020). Modeling for Predictors of Knowledge Score on Etiology and Prevention Strategies for Cervical Cancer Among Women of Reproductive Age in Ibadan. *JCO Global Oncology*. 2020; (6): 892.