



























- Mofolo A Innocent, Chigwenembe Maggie, Sichali S Dorothy, van der Horst M Charles.(2010). *Maternal or Infant Antiretroviral Drugs to Reduce HIV Transmission*. N Engl J Med, 362 (24):2271-81.
- [4]. Esther Ndonga, Zipporah Ng'ang'a, Erastus Muniu, Mohamed Karama and Martin Matu (2014) Barriers to Uptake and Effective Integration of PMTCT into SRH Services in Selected Health Facilities in Nairobi County, Kenya. *Journal of Pediatrics & Neonatal Care*. Volume 1(4).
- [5]. Fisher AA, Laing JE, Strocker J E (1998) *Handbook for family planning, operation research design in sampling*. Population Council. 40-45
- [6]. George Awungafac, Patrick Achiangia Njukeng, Juliana Ajoache Ndasi, and Lawrence Tanyi Mbuagbaw (2015) Prevention of mother-to-child transmission of the Human Immunodeficiency Virus: investigating the uptake and utilization of maternal and child health services in Tiko health district, Cameroon. *Pan Afr Med J*. 2015; 20: 20
- [7]. Gundel H, Katja S, Ilaria M, Chris B and Paulina M. (2005) *Analyzing awareness and Handbook for family planning, operation research design in sampling. Population Immunodeficiency virus; a study on acceptability by Nigerian women attending clinics. African Journal of Reproductive Health*.
- [8]. Health Bridge. (2007). *Challenges in the prevention of Mother to child transmission of HIV in Africa*.
- [9]. Katushabe J. (2006). *Knowledge and attitude of pregnant women on the use of prevention of mother to child transmission of HIV (PMTCT) services in Mbale regional hospital*.
- [10]. Moth I A, Ayayo ABCO, Kaseje DO (2000). *Assessment of utilization of PMTCT services at Nyanza Provincial Hospital, Kenya. Sahara J. PubMed*. 2005;2:244-50. *mother-to-child HIV transmission in resource-poor countries: Translating research into policy and practice. JAMA. PubMed*, 283:1175-1182.
- [11]. National AIDS and STI Control Programme (2009) *Sentinel Surveillance of HIV in Antenatal Clinics 2008/2009*. Ministry of Health, Nairobi, Kenya
- [12]. NASCOP, "Kenya National Guidelines for HIV Prevention and Management of People Who Use Drugs (PWUD)", Kenya. 2013.
- [13]. Perez, F., Mukotekwa, T, & Miller, A.(2004). "Implementing a rural programme of prevention of mother-to-child transmission of HIV in Zimbabwe: first 18 months of experience." *Tropical Medicine and International Health*, vol. 9, no. 7, pp. 774–783.
- [14]. Secka, E. (2010). *Men's involvement in care and support during pregnancy and childbirth: A qualitative study conducted in Gambia*.
- [15]. Semrau K, Kuhn L, Vwalika C, Kasonde P, Sinkala M, Kankasa C, Shutes E, Aldrovandi G, Thea DM (2005) Women in couples antenatal HIV counseling and testing are not more likely to report adverse social events. *AIDS*. 19(6): 603-9.
- [16]. Sirengo M, Muthoni L, Kellogg TA, Kim AA, Katana A (2014) Mother-to-child transmission of HIV in Kenya: results from a nationally representative study. *J Acquir Immune Defic Syndr* 66 Suppl 1S66–74.
- [17]. UNAIDS 2010. *Global Report. UNAIDS report on the global AIDS epidemic/2010*. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS (UNAIDS); 2010.
- [18]. UNAIDS (2013) *UNAIDS report on the global AIDS epidemic 2013*. Geneva
- [19]. UNAIDS AIDS (2001) *Declaration of Commitment on HIV/AIDS*. United General assembly. Geneva.
- [20]. UNICEF (2012) *Tracking Progress on child and maternal nutrition*. Assessed from [www.unicef.org/publications/files/Tracking\\_Progress\\_on\\_Child\\_and\\_Maternal\\_Nutrition\\_EN\\_110309.pdf](http://www.unicef.org/publications/files/Tracking_Progress_on_Child_and_Maternal_Nutrition_EN_110309.pdf), on 23<sup>rd</sup> February, 2015
- [21]. USAID; (2001): *Effective programs for preventing mother-To-Child HIV transmission*. Washington DC. The population Council: Washington, DC USA
- [22]. WHO (2013) *Reproductive Health and Research*. Assessed from <http://www.who.int/malaria/publications/atoz/iptp-sp-updated-policy-brief-24jan2014.pdf>, on 23<sup>rd</sup> February, 2017.
- [23]. Zolfo, M., Delvaux, T. & Tamburrini, E. (2005). *HIV/AIDS Knowledge and PMTCT Demand in Rural Zimbabwe*. Washington: BMJ Publishing group