















- [4]. Nigeria Demographic and Health Survey (NDHS). HIV/AIDS Survey, 2013. Available at <https://dhsprogram.com/pubs/pdf/FR293/FR293.pdf>. Accessed 19 March 2020.
- [5]. Fuge, T. G., Ayanto S. Y., and Gurmamo, F. L., 2015, Assessment of knowledge, attitude and practice about malaria and ITNs utilization among pregnant women in Shashogo District, Southern Ethiopia. *Malar J.* 2015, 14, 235.
- [6]. Ezire O., Adebayo S., Idogho, O., Bamgboye E., and Nwokolo E., 2015, Determinants of use of insecticide-treated nets among pregnant women in Nigeria *International Journal of Womens Health.* 7: 655–661.
- [7]. Adeniran, A, Goodman, O. O., Olatona, F. A., and Oluwole, E. O., 2016, Malaria prevention in pregnancy among traditional birth attendants in rural Lagos, Nigeria.
- [8]. Ifeoma, P., Okafor, Chinonso E., Esther, O., Olanrewaju, O., and Onigbogi, O., 2019, Malaria in pregnancy: A community-based study on the knowledge, perception, and prevention among Nigerian women. *J Family Med Prim Care* 8, 1359-64.
- [9]. Rogerson, S. J., Desai, M., Mayor, A., Sicuri, E., Taylor, S. M., Van, and Eijk A. M., 2018, Burden, pathology, and costs of malaria in pregnancy:
- [10]. Adaramola, E. F., and Babalola B. I., 2017, Factors influencing the utilization of insecticide treated nets among children under the age of five in Nigeria. Federal University Oye-Ekiti, Ekiti State Nigeria 2015. Available at: <http://uaps2015.princeton.edu/uploads/150778>. Accessed 14th Sep 2017.
- [11]. Inungu, J. N., Ankiba, N., Minelli, M., Mumford, V., Bolekela, D., and Mukoso, B., 2017, Use of insecticide-treated mosquito net among pregnant women and guardians of children under five in the Democratic Republic of the Congo. *Malar Res Treat*, 2017:1–10.
- [12]. Seacat, J. D., and Northrup, D., 2010, An information-motivation-behavioral skills assessment of curbside recycling behavior. *J Environ Psychol*, 30(4):393–401.
- [13]. Nganda, R. Y., Drakeley, C., Reyburn, H., and Marchant, T., 2004, Knowledge of malaria influences the use of insecticide treated nets but not intermittent presumptive treatment by pregnant women in Tanzania. *Malar J.* 3:42.
- [14]. Obol, J. H., Ononge, S., and Orach, C. G., 2013, Utilisation of insecticide treated nets among pregnant women in Gulu: a post conflict district in northern Uganda. *Afr Health Sci*, 13:962–9.
- [15]. Inungu, J. N., Ankiba, N., Minelli, M., Mumford, V., Bolekela, D., and Mukoso, B., 2017, Use of insecticide-treated mosquito net among pregnant women and guardians of children under five in the Democratic Republic of the Congo. *Malar Res Treat*, 2017:1–10.
- [16]. Balami, A. D., Salmiah S., Zulkefli, N A., Bachok, N., and Balami, E. L., 2019, An information-motivation-behavioural-skills analysis of long-lasting insecticidal net use among pregnant women in a hospital in North-Eastern Nigeria. *BMC Medical Research Methodology* 19:157.