













- [11]. Guiney, H. and L. Machado, Benefits of regular aerobic exercise for executive functioning in healthy populations. *Psychonomic bulletin & review*, 2013. 20(1): p. 73-86.
- [12]. Smith, P.J., et al., Aerobic exercise and neurocognitive performance: a meta-analytic review of randomized controlled trials. *Psychosomatic medicine*, 2010. 72(3): p. 239.
- [13]. Brown, A.D., et al., Effects of cardiorespiratory fitness and cerebral blood flow on cognitive outcomes in older women. *Neurobiology of Aging*, 2010. 31(12): p. 2047-2057.
- [14]. Endes K, Köchli S, Zahner L and Hanssen H (2019) Exercise and Arterial Modulation in Children: The EXAMIN YOUTH Study. *Front. Physiol.* 10:43. doi:10.3389/fphys.2019.00043.
- [15]. Ashmita Karki, Archana Shrestha, Narayan Subed (2019). Prevalence and associated factors of childhood overweight/obesity among primary school children in urban Nepal. *BMC Public Health*, 19:1055 <https://doi.org/10.1186/s12889-019-7406-9>.
- [16]. Hayman, L.L., et al., Cardiovascular Health Promotion in the Schools. *Circulation*, 2004. 110(15): p. 2266-2275.
- [17]. Hogstrom, G., A. Nordstrom, and P. Nordstrom, High aerobic fitness in late adolescence is associated with a reduced risk of myocardial infarction later in life: a nationwide cohort study in men. *Eur Heart J*, 2014. 35(44): p. 3133-40.
- [18]. Coimbra, S., et al., Physical exercise intervention at school improved hepcidin, inflammation, and iron metabolism in overweight and obese children and adolescents. *Pediatric research*, 2017. 82(5): p. 781-788.
- [19]. Kelley, G.A., K.S. Kelley, and R.R. Pate, Exercise and BMI z-score in Overweight and Obese Children and Adolescents: A Systematic Review and Network Meta-Analysis of Randomized Trials. *J Evid Based Med*, 2017. 10(2): p. 108-128.
- [20]. Wen, H. and L. Wang, Reducing effect of aerobic exercise on blood pressure of essential hypertensive patients: A meta-analysis. *Medicine (Baltimore)*, 2017. 96(11): p. e6150.
- [21]. Cornelissen, V.A., et al., Effects of aerobic training intensity on resting, exercise and post-exercise blood pressure, heart rate and heart-rate variability. *J Hum Hypertens*, 2010. 24(3): p. 175-82.
- [22]. Lemes, I.R., et al., Aerobic training reduces blood pressure and waist circumference and increases HDL-c in metabolic syndrome: a systematic review and meta-analysis of randomized controlled trials. *J Am Soc Hypertens*, 2018. 12(8): p. 580-588.
- [23]. Lee, K., et al., Effect of Aerobic and Resistance Exercise Intervention on Cardiovascular Disease Risk in Women With Early-Stage Breast Cancer: A Randomized Clinical Trial. *JAMA Oncol*, 2019. 5(5): p. 710-714.