





























limited setting: a randomized controlled trial of text message reminders. *AIDS*. 2011;25(6):825–34. pmid:21252632.

[30] Lester RT, Ritvo P, Mills EJ, Kariri A, Karanja S, Chung MH, *et al.* (2010). Effects of a mobile phone short message service on antiretroviral treatment adherence in Kenya (WelTel Kenya1): a randomized trial. *Lancet*. 2010;376(9755):1838–45. pmid:21071074 [doi:10.1016/S0140-6736(10)61997-6].

[31] Woods SP, Moran LM, Carey CL, Dawson MS, Iudicello JE, *et al.* (2008) Prospective memory in HIV infection: is “remembering to remember” a unique predictor of self-reported medication management? *Arch Clin Neuropsychology* 23:257–270.

[32] Chung MH, Richardson BA, Tapia K, Benki-Nugent S, Kiarie JN, Simoni JM, *et al.* (2011). A randomized controlled trial comparing the effects of counseling and alarm device on HAART adherence and virologic outcomes. *PLoS Medicine*. 2011;8(3): e1000422. pmid:21390262.

[33] Abdulrahman SA, Rampal L, Ibrahim F, Radhakrishnan AP, Kadir Shahar H, Othman N (2017) Mobile phone reminders and peer counseling improve adherence and treatment outcomes of patients on ART in Malaysia: A randomized clinical trial. *PLoS ONE* 12(5): e0177698. <https://doi.org/10.1371/journal.pone.0177698>.

[34] Kunutsor S, Walley J, Katabira E, Muchuro S, Balidawa H, Namagala E, *et al.* Using mobile phones to improve clinic attendance amongst an antiretroviral treatment cohort in rural Uganda: A cross-sectional and prospective study. *AIDS Behav*. 2010; 14(6): 1347±1353. <https://doi.org/10.1007/s10461-010-9780-2> PMID: 20700644.

[35] Sarna A, Luchters S, Geibel S, Chersich MF, Munyao P, Kaai S, *et al.* Short-and long-term efficacy of modified directly observed antiretroviral treatment in Mombasa, Kenya: a randomized trial. *J Acquir Immune Defic Syndr*. 2008; 48(5), 611±619. <https://doi.org/10.1097/QAI.0b013e3181806bf1> PMID:18645509.

[36] Rich ML, Miller AC, Niyigena P, Franke MF, Niyonzima JB, Soggi A, *et al.* (2012). Excellent Clinical Outcomes and High Retention in Care among Adults in a Community-Based HIV Treatment Program in Rural Rwanda *J Acquire Immune Defic Syndr*. 2012; 59(3): e35±e42. <https://doi.org/10.1097/QAI.0b013e31824476c4> PMID: 22156912.

[37] Mannheimer S, Friedland G, Matts J, Child C, Chesney M. The consistency of adherence to antiretroviral therapy predicts biologic outcomes for human immunodeficiency virus infected persons in clinical trials. *Clin Infect Dis*. 2002; 34:1115–21.

[38] El-Khatib Z, Ekstrom AM, Coovadia A, Abrams EJ, Petzold M, *et al.* (2011) Adherence and virologic suppression during the first 24 weeks on antiretroviral therapy among women in Johannesburg, South Africa - a prospective cohort study. *BMC Public Health* 11: 88.

[39] Sampaio-Sa M, Page-Shafer K, Bangsberg DR, *et al.* 100% adherence study: educational workshops vs video sessions to improve adherence among ART-naive patients in Salvador, Brazil. *AIDS Behaviour*. 2008;12(4 Suppl): S54–S62.

[40] Garcia R, Ponde M, Lima M, Souza AR, Stolze SM, Badaro R. Lack of effect of motivation on the adherence of HIV-positive/AIDS patients to antiretroviral treatment. *Braz J Infect Dis*. 2005;9(6):494–499.