

- Development of infrastructure and description of study site. *Am. J. Trop. Med. Hyg.* (2003);68(4 suppl):3-9.
- [25]Ter Kuile FO, Terloun DJ, Kariuki SK, et al. Impact of Permethrin Treated Bed Nets on Malaria, Anemia and Growth in Infants in an Area of Intense Perennial Malaria Transmission in Western Kenya. *Am. J. Trop. Med. Hyg.* (2003): 68(4 suppl): 68-77.
- [26]Gamble C, Ekwaru JP, Ter Kuile FO. Insecticide- treated nets for preventing malaria in pregnancy. *Cochrane Database of Syst Rev.* (2006) Apr 19; (2): CD003755.
- [27]Negash K, Jima D, Nafo-Traore F, et al. Ethiopian Roll Back Malaria Consultative Mission: Essential Actions to Support the Attainment of the Abuja Targets. 16th to 20th February (2004).
- [28]Anyanwu EC, Ehiri JE, Kanu I, Morad M, Ventegodt S, Merrick J. Assessing the effects of long-term exposure to insecticide treated mosquito nets in the control of malaria in endemic regions. *Scientific world Journal* (2004); 4:978-88.
- [29]Roll Back Malaria. (2001-2010) United Nations Decade to Roll Back Malaria: Insecticide-treated Mosquito Nets. RBM info sheet. Accessed through: <http://www.rbm.who.int>. (Accessed on: 15/ 08/ 2006).
- [30]Abuja Declaration and the Plan of Action: An Extract from the African Summit on Roll Back Malaria, Abuja, 25 April (2000) (WHO/CDS/RBM/ 2000.17). Available at: http://www.rbm.who.int/docs/abuja_declaration_final.htm. (Accessed on: 15/ 08/ 2006).
- [31]Worrall E, Hill J, Webster J, and Mortimer J. Experience of targeting subsidies on insecticide-treated nets: What do we know and what are the knowledge gaps? *Trop. Med. Int. Health* (2005); 10(1): 19-31.
- [32]Hill J, Lines J, Rowland M. Insecticide-treated nets. *Adv Parasitol.* (2006); 61:77-128.
- [33]Net Mark research (1999-2006). Available at: <http://www.netmarkafrica.org/research>. (Accessed on: 15/ 08/ 2006).
- [34]Macintyre K, Keating J, Okbaldt YB, et al. Rolling out insecticide treated nets in Eritrea: examining the determinants of possession and use in malarious zones during rainy seasons. *Trop Med Int Health.* (2006) Jun; 11(6): 824-33.
- [35]Tsuyuoko R, Midizi SM, Dziva P, Makunike B. The acceptability of insecticide treated mosquito nets among community members in Zimbabwe. *Cent Afr J Med.* (2002) Jul-Aug; 48 (7-8):87-91.
- [36]Morgan DW and Krejcie, RV. (1970). Determining Sample size for research activities of Minnesota: USA.
- [37]Deressa, W., Ali, A., Enquosellassie F. Knowledge, attitude and practice about malaria, the mosquito and antimalarial drugs in a rural community. *Ethiop J Health Dev* (2003); 17(2): 99-104.
- [38]Yeneneh, H., Gyorkos, T.W., Joseph, L., Pickering, J. and Tedla, S. Antimalarial drug utilization by women in Ethiopia: a knowledge-attitudes-practice study. *Bull World Health Org* (1993); 71: 763-772.
- [39]Ongore, D., Kamunvi, F., Knight, R., and Minawa, A. A study of knowledge, attitudes and practices (KAP) of a rural community on malaria and the mosquito vector. *East Afr Med J* (1989); 66(2):79-89.
- [40]Klein, R.E., Weller, S.C., Zeissig, R., Richards, F.O. and Ruebush, T.K. Knowledge, beliefs, and practices in relation to malaria transmission and vector control in Guatemala. *Am J Trop Med Hyg* (1995); 52(5): 383-388.
- [41]Gebreyesus, T.A., Witten, K.H., Getachew, A. et al. The community-based malaria control in Tigray, northern Ethiopia. A review of programme set-up, activities, outcomes and impact. *Parasitologia* (2000); 42: 255-240.
- [42]Kidane, G. and Morrow, R.H. Teaching mothers to provide home treatment of malaria in Tigray, Ethiopia: a randomized trial. *Lancet* (2000); 356(9229): 550-555.
- [43]Binka, F.N. and Adonso. P. Acceptability and use of insecticide impregnated bed nets in northern Ghana. *Trop Med Int Health* (1997); 2(5): 499-507.