















Buchsbaum S, Moeller G, Hay Burgess DC, 2006. Requirements for high impact diagnostics in the developing world. *Nature*. 2006 Nov 23; 444 Suppl 1():73-9. Date of Access: 29/06/2020. <https://www.nature.com/articles/nature05448>.

[11] Burgess DCH, Wasserman J, Dahl CA, Girosi F, Olmsted SS, Keeler EB, et al. 2007. Estimating the Global Health Impact of Improved Diagnostic Tools for the Developing World. Santa Monica, CA: RAND Corporation; 2007. Date of Access: 29/06/2020. [http://www.rand.org/pubs/research\\_briefs/RB9293.html](http://www.rand.org/pubs/research_briefs/RB9293.html).

[12] Jani IV, Meggi B, Loquiha O, *et al.* Effect of point-of-care early infant diagnosis on antiretroviral therapy initiation and retention of patients. *AIDS*. 2018;32(11):1453-1463.

doi:10.1097/QAD.0000000000001846.

[13] Flavia Bianchi, Jennifer Cohn, Emma Sacks, Rebecca Bailey, Jean-Francois Lemaire and Rhoderick Machekano. 2019. Evaluation of a routine

point-of-care intervention for early infant diagnosis of HIV: an observational study in eight African countries. *The Lancet HIV* Volume 6, Issue 6, June 2019, Pages e373-e38. Date Accessed: 15/5/2020. [https://doi.org/10.1016/S2352-3018\(19\)30033-5](https://doi.org/10.1016/S2352-3018(19)30033-5).

[14] Pharr, Jennifer R., Obiefune, Michael C.; Ezeanolue, Chinenye O.; Osuji, Alice; Ogidi, Amaka G.; Gbadamosi, Semiu, Patel, Dina, Iwelunmor, Juliet; Yang, Wei; Ogedegbe, Gbenga; Ehiri, John E.; Sam-Agudu, Nadia A.; Ezeanolue, Echezona E. 2016. Linkage to Care, Early Infant Diagnosis, and Perinatal Transmission Among Infants Born to HIV-Infected Nigerian Mothers: Evidence from the Healthy Beginning Initiative, *JAIDS Journal of Acquired Immune Deficiency Syndromes*: August 1, 2016 - Volume 72 - Issue - p S154-S160. Date of Access: 5/04/2020.

<https://doi.org/10.1097/QAI.0000000000001051>.